2025

Signature of Student

PERTH AMBOY PUBLIC SCHOOLS

2026

Athletic Department Perth Amboy, NJ 08861 (732) 376-6230 Ext 40-407 / 40-408

Athletic Consent for:		Sport(s)			
		. ,	·			
Name of Student (Print)			•	Gra	ade	ID#
Address						
Home Phone	Work Phone				Cell Phone	
Emergency Contact Name	 Er	nergenc	y Co	ontact	Phone Numbe	r
l am enrolled in the Perth Amb	ooy School District	YES	1	NO	(Circle appro	priate response)
l am a transfer student		YES	1	NO	(Circle appro	priate response)
If a transfer student - I have competed in thes	se sports at my prior o	district:			(List Sports T	eam(s) Here)
hereby give consent for my child	ussion, Covid19, NJSIAA 5 days of the start of properties Ier should fill out the Epother The Athletic Trainer/Nurblete and submit the form The Adelay in a delay in	ractice for pipen E se at leasorms as partici	or th mei ast <u>t</u> ins pati	ne spo rgenc hirty (structe	ort in which party Care Plan of 30) days prior ed thirty (30) the tryouts.	rticipation is desired. or the Asthma Action Plan to tryouts.
 Represent Perth Amboy Public Schools in Accompany any school team in which yo 						f-town trips.
authorize the school to provide any emergency ravel.	medical care that may	become	ne	cessa	ry for my child	during such activities or
Warning: such activities involve the potential for of protective equipment, and strict observance of severe as to result in paralysis, total disability or conderstand this warning.	rules; injuries are still a	a possib	ility.	On r	are occasions,	, these injuries can be so
authorize the release of academic information Coaching Staff; I guarantee that the above inf regarding my child's participation, Code of Et	ormation is correct a	nd that	l ag	ree to	abide by all	the above information
Signature of Parent/Guardian	and the control of th	<u>_</u>			Date	
By signing this consent, I acknowledge that I information stated above; conduct myself acc equipment issued to me.						=

Date

This form should be maintained by the healthcare provider completing the physical exam (medical home). It should not be shared with schools. The medical eligibility form is the only form that should be submitted to a school. The physical exam must be completed by a healthcare provider who is a licensed physician, advanced practice nurse or physician assistant who has completed the Student-Athlete Cardiac Assessment Professional Development module hosted by the New Jersey Department of Education.

■ PREPARTICIPATION PHYSICAL EVALUATION (Interim Guidance) HISTORY FORM

Note: Complete and sign this form (with your parents	-	•	*				
Name:	Sport(s)	D	tie of birm:				
Sex assigned at birth (F, M, or intersex):	•			ner gender):			
Have you had COVID-19? (check one): □ Y □ N	1						
Have you been immunized for COVID-19? (check o	one): □Y □N		u had: □ One shot [□ Booster date(s)				
List past and current medical conditions.							
Have you ever had surgery? If yes, list all past surgic	al procedures						
Medicines and supplements: List all current prescrip	tions, over-the-co	unter medicines, a	nd supplements (herbal	and nutritional).			
Do you have any allergies? If yes, please list all you	r allergies (ie, me	dicines, pollens, fo	ood, stinging insects).				
Patient Health Questionnaire Version 4 (PHQ-4) Over the last 2 weeks, how often have you been bothered by any of the following problems? (Circle response.)							
		Several days	Over half the days	Nearly every day			
Feeling nervous, anxious, or on edge	0	1	2	3			
Not being able to stop or control worrying	0	1	2	3 '			
Little interest or pleasure in doing things	0	1	2	3			
Feeling down, depressed, or hopeless	0	1	2	3			
(A sum of ≥3 is considered positive on either s	ubscale [question	s 1 and 2, or ques	tions 3 and 4] for scree	ening purposes.)			
SEVERAL OUTSTONE							

(Exp	IERAL QUESTIONS lain "Yes" answers at the end of this form. Circle stions if you don't know the answer.)	Yes	No
1.	Do you have any concerns that you would like to discuss with your provider?		
2.	Has a provider ever denied or restricted your participation in sports for any reason?		
3.	Do you have any ongoing medical issues or recent illness?		
HEA	RT HEALTH QUESTIONS ABOUT YOU	Yes	No
4.	Have you ever passed out or nearly passed out during or after exercise?		
5.	Have you ever had discomfort, pain, tightness, or pressure in your chest during exercise?		
6.	Does your heart ever race, flutter in your chest, or skip beats (irregular beats) during exercise?		
7.	Has a doctor ever told you that you have any heart problems?		
8.	Has a doctor ever requested a test for your heart? For example, electrocardiography (ECG) or echocardiography.		

	ART HEALTH QUESTIONS ABOUT YOU ONTINUED)		Yes	Na
9	Do you get light-headed or feel shorter of breathan your friends during exercise?	ath		
10	. Have you ever had a seizure?			
HEA	RT HEALTH QUESTIONS ABOUT YOUR FAMILY	Unsure	Yes	No
11.	Has any family member or relative died of heart problems or had an unexpected or unexplained sudden death before age 35 years (including drowning or unexplained car crash)?			
12.	Does anyone in your family have a genetic heart problem such as hypertrophic cardiomyopathy (HCM), Marfan syndrome, arrhythmogenic right ventricular cardiomyopathy (ARVC), long QT syndrome (LQTS), short QT syndrome (SQTS), Brugada syndrome, or catecholaminergic polymorphic ventricular tachycardia (CPVT)?			
13.	Has anyone in your family had a pacemaker or an implanted defibrillator before age 35?			

.(e)	NE AND JOINT QUESTIONS	Yes	No	ME	DICAL QUESTIONS (CONTINUED)	(es
14.	Have you ever had a stress fracture or an injury to a			25	Do you worry about your weight?	
	bone, muscle, ligament, joint, or tendon that caused you to miss a practice or game?			26	Are you trying to or has anyone recommended that you gain or lose weight?	
15.	Do you have a bone, muscle, ligament, or joint injury that bothers you?			27	Are you on a special diet or do you avoid certain types of foods or food groups?	
ME	DICAL QUESTIONS	Yes	No	28	Have you ever had an eating disorder?	
16.	Do you cough, wheeze, or have difficulty breathing during or after exercise?				STRUAL QUESTIONS N/A Y Have you ever had a menstrual period?	es
17.	Are you missing a kidney, an eye, a testicle, your spleen, or any other organ?				How old were you when you had your first menstrual period?	
18.	Do you have groin or testicle pain or a painful bulge			31.	When was your most recent menstrual period?	
	or hernia in the groin area?	<u> </u>		32.	How many periods have you had in the past 12	
19.	Do you have any recurring skin rashes or rashes that come and go, including herpes or methicillin-resistant Staphylococcus aureus (MRSA)?			Exp!	months? sin "Yes" answers here.	
20.	Have you had a concussion or head injury that caused confusion, a prolonged headache, or memory problems?					
21.	Have you ever had numbness, had tingling, had weakness in your arms or legs, or been unable to move your arms or legs after being hit or falling?			***************************************		
22.	Have you ever become ill while exercising in the heat?			***************************************		
23.	Do you or does someone in your family have sickle cell trait or disease?					
24.	Have you ever had or do you have any problems with your eyes or vision?					

Signature of parent or guardian: ____

^{© 2023} American Academy of Family Physicians, American Academy of Pediatrics, American College of Sports Medicine, American Medical Society for Sports Medicine, American Orthopaedic Society for Sports Medicine, and American Osteopathic Academy of Sports Medicine. Permission is granted to reprint for noncommercial, educational purposes with acknowledgment.

This form should be maintained by the healthcare provider completing the physical exam (medical home). It should not be shared with schools. The Medical Eligibility Form is the only form that should be submitted to a school.

PREPARTICIPATION PHYSICAL EVALUATION ATHLETES WITH DISABILITIES FORM: SUPPLEMENT TO THE ATHLETE HISTORY

Name:Date of birth:		
1. Type of disability:		
2. Date of disability:		
3. Classification (if available):		
4. Cause of disability (birth, disease, injury, or other):		
5. List the sports you are playing:		
3. List the sports you are playing.		
6. Do you regularly use a brace, an assistive device, or a prosthetic device for daily activities?	Yes	No
7. Do you use any special brace or assistive device for sports?		
8. Do you have any rashes, pressure sores, or other skin problems?		
9. Do you have a hearing loss? Do you use a hearing aid?		
10. Do you have a visual impairment? 11. Do you use any special devices for bowel or bladder function?		
12. Do you have burning or discomfort when urinating? 13. Have you had autonomic dysreflexia?		
14. Have you ever been diagnosed as having a heat-related (hyperthermia) or cold-related (hypothermia) illness?		
15. Do you have muscle spasticity?		
16. Do you have frequent seizures that cannot be controlled by medication?		
Explain "Yes" answers here.		
	7.11	
Please indicate whether you have ever had any of the following conditions:		
. to a continuous which is you have ever had any or the following conditions.	I w. I	, vie
	Yes	No
	1 1	1600010250000
Atlantoaxial instability Radiographic (x-ray) evaluation for atlantoaxial instability		
Radiographic (x-ray) evaluation for atlantoaxial instability		
Radiographic (x-ray) evaluation for atlantoaxial instability Dislocated joints (more than one)		
Radiographic (x-ray) evaluation for atlantoaxial instability Dislocated joints (more than one) Easy bleeding		
Radiographic (x-ray) evaluation for atlantoaxial instability Dislocated joints (more than one) Easy bleeding Enlarged spleen		
Radiographic (x-ray) evaluation for atlantoaxial instability Dislocated joints (more than one) Easy bleeding Enlarged spleen Hepatitis		
Radiographic (x-ray) evaluation for atlantoaxial instability Dislocated joints (more than one) Easy bleeding Enlarged spleen Hepatitis Osteopenia or osteoporosis		
Radiographic (x-ray) evaluation for atlantoaxial instability Dislocated joints (more than one) Easy bleeding Enlarged spleen Hepatitis Osteopenia or osteoporosis Difficulty controlling bowel		
Radiographic (x-ray) evaluation for atlantoaxial instability Dislocated joints (more than one) Easy bleeding Enlarged spleen Hepatitis Osteopenia or osteoporosis Difficulty controlling bowel Difficulty controlling bladder		
Radiographic (x-ray) evaluation for atlantoaxial instability Dislocated joints (more than one) Easy bleeding Enlarged spleen Hepatitis Osteopenia or osteoporosis Difficulty controlling bowel Difficulty controlling bladder Numbness or tingling in arms or hands		
Radiographic (x-ray) evaluation for atlantoaxial instability Dislocated joints (more than one) Easy bleeding Enlarged spleen Hepatitis Osteopenia or osteoporosis Difficulty controlling bowel Difficulty controlling bladder Numbness or tingling in arms or hands Numbness or tingling in legs or feet		
Radiographic (x-ray) evaluation for atlantoaxial instability Dislocated joints (more than one) Easy bleeding Enlarged spleen Hepatitis Osteopenia or osteoporosis Difficulty controlling bowel Difficulty controlling bladder Numbness or tingling in arms or hands Numbness or tingling in legs or feet Weakness in arms or hands		
Radiographic (x-ray) evaluation for atlantoaxial instability Dislocated joints (more than one) Easy bleeding Enlarged spleen Hepatitis Osteopenia or osteoporosis Difficulty controlling bowel Difficulty controlling bladder Numbness or tingling in arms or hands Numbness or tingling in legs or feet Weakness in arms or hands Weakness in legs or feet		
Radiographic (x-ray) evaluation for atlantoaxial instability Dislocated joints (more than one) Easy bleeding Enlarged spleen Hepatitis Osteopenia or osteoporosis Difficulty controlling bowel Difficulty controlling bladder Numbness or tingling in arms or hands Numbness or tingling in legs or feet Weakness in arms or hands Weakness in legs or feet Recent change in coordination		
Radiographic (x-ray) evaluation for atlantoaxial instability Dislocated joints (more than one) Easy bleeding Enlarged spleen Hepatitis Osteopenia or osteoporosis Difficulty controlling bowel Difficulty controlling bladder Numbness or tingling in arms or hands Numbness or tingling in legs or feet Weakness in arms or hands Weakness in legs or feet Recent change in coordination Recent change in ability to walk		
Radiographic (x-ray) evaluation for atlantoaxial instability Dislocated joints (more than one) Easy bleeding Enlarged spleen Hepatitis Osteopenia or osteoporosis Difficulty controlling bowel Difficulty controlling bladder Numbness or tingling in arms or hands Numbness or tingling in legs or feet Weakness in arms or hands Weakness in legs or feet Recent change in coordination Recent change in ability to walk Spina bifida		
Radiographic (x-ray) evaluation for atlantoaxial instability Dislocated joints (more than one) Easy bleeding Enlarged spleen Hepatitis Osteopenia or osteoporosis Difficulty controlling bowel Difficulty controlling bladder Numbness or tingling in arms or hands Numbness or tingling in legs or feet Weakness in arms or hands Weakness in legs or feet Recent change in coordination Recent change in ability to walk Spina bifida Latex allergy		
Radiographic (x-ray) evaluation for atlantoaxial instability Dislocated joints (more than one) Easy bleeding Enlarged spleen Hepatitis Osteopenia or osteoporosis Difficulty controlling bowel Difficulty controlling bladder Numbness or tingling in arms or hands Numbness or tingling in legs or feet Weakness in arms or hands Weakness in legs or feet Recent change in coordination Recent change in ability to walk Spina bifida		
Radiographic (x-ray) evaluation for atlantoaxial instability Dislocated joints (more than one) Easy bleeding Enlarged spleen Hepatitis Osteopenia or osteoporosis Difficulty controlling bowel Difficulty controlling bladder Numbness or tingling in arms or hands Numbness or tingling in legs or feet Weakness in arms or hands Weakness in legs or feet Recent change in coordination Recent change in ability to walk Spina bifida Latex allergy		
Radiographic (x-ray) evaluation for atlantoaxial instability Dislocated joints (more than one) Easy bleeding Enlarged spleen Hepatitis Osteopenia or osteoporosis Difficulty controlling bowel Difficulty controlling bladder Numbness or tingling in arms or hands Numbness or tingling in legs or feet Weakness in arms or hands Weakness in arms or hands Weakness in legs or feet Recent change in coordination Recent change in ability to walk Spina bifida Latex allergy Explain "Yes" answers here.	ete and correct	
Radiographic (x-ray) evaluation for atlantoaxial instability Dislocated joints (more than one) Easy bleeding Enlarged spleen Hepatitis Osteopenia or osteoporosis Difficulty controlling bowel Difficulty controlling bladder Numbness or tingling in arms or hands Numbness or tingling in legs or feet Weakness in arms or hands Weakness in legs or feet Recent change in coordination Recent change in ability to walk Spina bifida Latex allergy	ete and correct	
Radiographic (x-ray) evaluation for atlantoaxial instability Dislocated joints (more than one) Easy bleeding Enlarged spleen Hepatitis Osteopenia or osteoporosis Difficulty controlling bowel Difficulty controlling bladder Numbness or tingling in arms or hands Numbness or tingling in legs or feet Weakness in arms or hands Weakness in arms or hands Weakness in legs or feet Recent change in coordination Recent change in ability to walk Spina bifida Latex allergy Explain "Yes" answers here. hereby state that, to the best of my knowledge, my answers to the questions on this form are completed.	ete and correct	

^{© 2019} American Academy of Family Physicians, American Academy of Pediatrics, American College of Sports Medicine, American Medicine, American Medicine, American Medicine, American Medicine, American Medicine, and American Osteopathic Academy of Sports Medicine. Permission is granted to reprint for roncommercial, educational purposes with acknowledgment.

This form should be maintained by the healthcare provider completing the physical exam (medical home). It should not be shared with schools. The medical eligibility form is the only form that should be submitted to a school. The physical exam must be completed by a healthcare provider who is a licensed physician, advanced practice nurse or physician assistant who has completed the Student - Athlete Cardiac Assessment Professional Development module Hosted by the New Jersey Department of Education.

PREPARTICIPATION PHYSICAL EVALUATION (Interim Guidance) PHYSICAL EXAMINATION FORM

N B N 4 W/R WIN WIN WINDER B N H H H H H W H W N W H W N W H W N W H W N W H W N W H W N W H W N W H W N W W H W N W W H W N W W N W N			
Name:	Date	of birth:	
 PHYSICIAN REMINDERS 1. Consider additional questions on more-sensitive issues. • Do you feel stressed out or under a lot of pressure? • Do you ever feel sad, hopeless, depressed, or anxious? • Do you feel safe at your home or residence? • Have you ever tried cigarettes, e-cigarettes, chewing tobacco, snuff, or dip? • During the past 30 days, did you use chewing tobacco, snuff, or dip? • Do you drink alcohol or use any other drugs? • Have you ever taken anabolic steroids or used any other performance-enh • Have you ever taken any supplements to help you gain or lose weight or in • Do you wear a seat belt, use a helmet, and use condoms? 2. Consider reviewing questions on cardiovascular symptoms (Q4–Q13 of History 	nancing supplement? mprove your performa	ance?	
EXAMINATION			
Height: Weight:			
BP: / (/) Pulse: Vision: R 20/ COVID-19 VACCINE Previously received COVID-19 vaccine: □ Y □ N	L 20/	Corrected: UY	□N
Administered COVID-19 vaccine at this visit: \(\text{Y} \) \(\text{N} \) If yes: \(\text{First dose} \)	□ Second does □ II	aird does - Roos	tor datale)
Administered COVID-19 vaccine at this visit:	9	NORMAL	
Eyes, ears, nose, and throat Pupils equal Hearing	The second secon		
Lymph nodes			
Heart [®] Murmurs (auscultation standing, auscultation supine, and ± Valsalva maneuver	•)		
Lungs			
Abdomen			
Skin • Herpes simplex virus (HSV), lesions suggestive of methicillin-resistant Staphyloc tinea corporis	occus aureus (MRSA)	l, or	
Neurological			
MUSCULOSKELETAL Neck		NORMAL	ABNORMAL FINDINGS
Back			
Shoulder and arm	A Administration of the Control of t		
Elbow and forearm			,
Wrist, hand, and fingers			
Hip and thigh			
Knee			
Leg and ankle			
Foot and toes			
Functional			
Double-leg squat test, single-leg squat test, and box drop or step drop test			
Consider electrocardiography (ECG), echocardiography, referral to a cardiologist nation of those. Jame of health care professional (print or type):		Da	nation findings, or a combi- te:
address:			MD DO ND D

© 2019 American Academy of Family Physicians, American Academy of Pediatrics, American College of Sports Medicine, American Medical Society for Sports Medicine, American Orthopaedic Society for Sports Medicine, and American Osteopathic Academy of Sports Medicine. Permission is granted to reprint for noncommercial, educational purposes with acknowledgment.

Preparticipation Physical Evaluation Medical Eligibility Form

The Medical Eligibility Form is the only form that should be submitted to school. It should be kept on file with the student's school health record.

Student	Athlete's Name	Date of	Birth	
Date of	Exam			
0	Medically eligible for all sports without restriction			
0	Medically eligible for all sports without restriction with re	ecommendatio	ns for further evaluation or treatment of	
o	Medically eligible for certain sports			
0	Not medically eligible pending further evaluation			
0	Not medically eligible for any sports			
Recomm	nendations:			
athlete de the physicondition	viewed the history form and examined the student named of ones not have apparent clinical contraindications to practice ical examination findings- are on record in my office and constant arise after the athlete has been cleared for participation, and the potential consequences are completely explained to	and can partical and be made averaged the made averaged the physician in t	sipate in the sport(s) as outlined on this form. A copy allable to the school at the request of the parents. If may rescind the medical eligibility until the problem is	of
Signature	e of physician, APN, PA		Office stamp (optional)	
Address:				
Name of	healthcare professional (print)			
l certify l Education	I have completed the Cardiac Assessment Professional Dev	elopment Mo	dule developed by the New Jersey Department of	
Signature	e of healthcare provider			
	Shared Hea	lth Informati	on	
Allergies				
Medicatio	ons:			
Other infor	mation:			
mergency	Contacts:			
merican Or	rican Academy of Family Physicians, American Academy of Pediatrics, Ame thopaedic Society for Sports Medicine, and American Osteopathic Academy tth acknowledgment.	of Sports Medicine	ports Medicine, American Medical Society for Sports Medicine, Permission is granted to reprint for noncommercial, educational School Physician:	
	*This form has been modified to meet the statutes set forth by No	ew Jersey.	Reviewed on	ate)
			(D	a10)

(School Physician Signature)

PERTH AMBOY PUBLIC SCHOOLS

ATHLETIC DEPARTMENT

931 Convery Blvd Perth Amboy, NJ 08861

Use and Misuse of Opioid Drugs Fact Sheet

Student-Athlete and Parent/Guardian Sign-Off

In accordance with N.J.S.A. 18A:40-41.10, public school districts, approved private schools for students with disabilities, and nonpublic schools participating in an interscholastic sports program must distribute this Opioid Use and Misuse Educational Fact Sheet to all student-athletes and cheerleaders. In addition, schools and districts must obtain a signed acknowledgement of receipt of the fact sheet from each student-athlete and cheerleader, and for students under age 18, the parent or guardian must also sign.

This sign-off sheet is due to the appropriate school personnel as determined by your district prior to the first official practice session of the spring 2018 athletic season (March 2, 2018, as determined by the New Jersey State Interscholastic Athletic Association) and annually thereafter prior to the student-athlete's or cheerleader's first official practice of the school year. Opioid Video is located at: https://youtu.be/3Rz6rkwpAx8

NJSIAA OPIOID POLICY ACKNOWLEDGEMENT

We have viewed the NJ CARES educational video on the risks of opioid use for high school athletes. We understand the NJSIAA policy that requires students, and their parents(s)/guardian(s) if a student is under the age of 18, to view this video and sign this acknowledgement.

o view this video	and sign this acknowledgement.	
	Name of School:	Perth Amboy High School
	Name of School District:	Perth Amboy Public Schools
/We acknowledge	e that we received and reviewed	the Educational Fact Sheet on the Use and Misuse of Opioid Drugs.
itudent ID#	Stud	dent Signature:
arent/Guardian s	Signature:	Date:

State of New Jersey DEPARTMENT OF EDUCATION

Sudden Cardiac Death Pamphlet Sign-Off Sheet

Name of School District: Name of Local School:	Perth Amboy Public Schools Perth Amboy High School	
I/We acknowledge that we repaired	eceived and reviewed the Sudden Cardiac Death in Young Athletes	
Student Signature:	Student ID#	-
Parent or Guardian Signati	ire:	
Date:	·	



NJSIAA STEROID TESTING POLICY

CONSENT TO RANDOM TESTING

In Executive Order 72, issued December 20, 2005, Governor Richard Codey directed the New Jersey Department of Education to work in conjunction with the New Jersey State Interscholastic Athletic Association (NJSIAA) to develop and implement a program of random testing for steroids, of teams and individuals qualifying for championship games.

Beginning in the Fall 2006 sports season, any student-athlete who possesses, distributes, ingests or otherwise uses any of the banned substances on the attached page, without written prescription by a fully-licensed physician, as recognized by the American Medical Association, to treat a medical condition, violates the NJSIAA's sportsmanship rule, and is subject to NJSIAA penalties, including ineligibility from competition.

Athlete may submit supplement and medication to Drug Free Sport AXISTM to receive information regarding banned substances or safety issue. Athlete or parents may login to the NJSIAA account at <u>axis.drugfreesport.com</u> using the password **njsports**.

The NJSIAA will test certain randomly selected individuals and teams that qualify for a state championship tournament or state championship competition for banned substances. The results of all tests shall be considered confidential and shall only be disclosed to the student, his or her parents and his or her school. No student may participate in NJSIAA competition unless the student and the student's parent/guardian consent to random testing.

By signing below, we consent to random testing in accordance with the NJSIAA steroid testing policy. We understand that, if the student or the student's team qualifies for a state championship tournament or state championship competition, the student may be subject to testing for banned substances.

Signature of Student-Athlete	Print Student-Athlete's Name	Date
Signature of Parent/Guardian	Print Parent/Guardian's Name	Date

necessary.

• To recover, cognitive rest is just as important as physical rest. Reading, texting, computer use and even watching movies can slow down recovery. Limit screen time during recovery.

Students who have sustained a concussion may not return to practice or competition until they receive written clearance from a physician trained in the evaluation and management of concussion and complete the graduated <u>Six-step return to play protocol outlined</u> by the <u>CDC</u>:

Step 1: Back to regular activities (such as school)

Athletes or cheerleaders are back to their regular activities (such as school).

Step 2: Light aerobic activity

Begin with light aerobic exercise only to increase an athlete's heart rate. This means about 5 to 10 minutes on an exercise bike, walking, or light jogging. No weightlifting at this point.

Step 3: Moderate activity

Continue with activities to increase an athlete's heart rate with body or head movement. This includes moderate jogging, brief running, moderate-intensity stationary biking, moderate-intensity weightlifting (less time and/or less weight from their typical routine).

Step 4: Heavy, non-contact activity

Add heavy non-contact physical activity, such as sprinting/running, high-intensity stationary biking, regular weightlifting routine, non-contact sport-specific drills (in 3 planes of movement).

Step 5: Practice & full contact

Athletes may return to practice and full contact (if appropriate for the sport) in controlled practice.

Step 6: Competition

Young athletes may return to competition.

For further information on Sports-Related Concussions and other Head Injuries, please visit:

- CDC Heads Up
- Keeping Heads Healthy

Student athlete's name (print)	Student athlete's signature	Date	
Parent / Guardian name (print)	Parent / Guardian signature	Date	



Sports-Related Concussion and Head Injury Fact Sheet and Parent/Guardian Acknowledgement Form

A concussion is a traumatic brain injury that can be caused by a blow to the head or body that disrupts the normal functioning of the brain. This sudden movement can cause the brain to bounce around or twist in the skull, creating chemical changes in the brain and sometimes stretching and damaging brain cells, disrupting the way the brain normally functions. Concussions can cause significant and sustained neuropsychological impairment affecting balance, reading (tracking), problem solving, planning, memory, attention, concentration, and behavior. Concussions can range from mild to severe. Having a concussion increases the risk of sustaining another concussion. Second-impact syndrome may occur when a person sustains a second concussion while still experiencing symptoms of a previous concussion. It can lead to severe impairment and even death.

Requirements addressing sports-related concussions and head injuries for student athletes and cheerleaders

- All school districts, charter, and non-public schools that participate in interscholastic sports are required to distribute this educational fact to all student athletes and cheerleaders and obtain a signed acknowledgment from each parent/guardian and student-athlete.
- Each school district, charter, and non-public school shall develop a written policy describing the
 prevention and treatment of sports-related concussion and other head injuries sustained by
 interscholastic student-athletes and cheerleaders.
- Any cheerleader or student-athlete who participates in an interscholastic sports program and is suspected of sustaining a concussion will be immediately removed from competition or practice. The student-athlete will not be allowed to return to competition or practice until they have written clearance from a physician trained in concussion treatment and have completed his/her district's graduated return-to-play protocol.

Quick Facts

- Most concussions do not involve loss of consciousness.
- You can sustain a concussion even if you do not hit your head.
- A blow elsewhere on the body can transmit an "impulsive" force to the brain and cause a concussion.
- Signs and symptoms of concussion can show up right after an injury or may not appear or be noticed until hours or days after the injury.

Signs of Concussions (Observed by Coach, Athletic Trainer, Parent/Guardian/Caregiver, Teammate, and others)

- Appears dazed or stunned
- Forgets plays or demonstrates short term memory difficulties (e.g., unsure of game, opponent)
- Exhibits difficulties with balance, coordination, concentration, and attention

- Answers questions slowly or inaccurately
- Is unable to recall events prior to or after the hit or fall

Symptoms of Concussion (Reported by Student-Athlete)

- Headache
- Nausea/vomiting
- Balance problems or dizziness
- Double vision or changes in vision trouble reading
- · Sensitivity to light/sound
- Feeling of sluggishness or fogginess fatigue
- Difficulty with concentration, short term memory, and/or confusion

Dangerous Signs & Symptoms of a Concussion

- New onset of symptoms
- One pupil is larger than the other
- Drowsiness or inability to wake up
- A headache that gets worse and does not go away
- Slurred speech, weakness, numbness, or decreased coordination
- Repeated vomiting, nausea, or seizures (shaking or twitching)
- Unusual behavior, increased confusion, restlessness, or agitation
- Loss of consciousness (passed out/knocked out); even a brief loss of consciousness should be taken seriously.

What should a student-athlete do if they think they have a concussion?

- Do not hide it. Tell your athletic trainer, coach, school nurse, or parent/guardian.
- Report it. Do not return to competition or practice with symptoms of a concussion or head injury.
- Take time to recover. If you have a concussion, your brain needs time to heal. While your brain is healing you are much more likely to sustain a second concussion.

What can happen if a student-athlete continues to play with a concussion or returns to play to soon?

- Continuing to play with the signs and symptoms of a concussion leaves the student-athlete vulnerable to second impact syndrome.
- Second impact syndrome is when a student-athlete sustains a second concussion while still having symptoms from a previous concussion or head injury.
- Second impact syndrome can lead to severe impairment and even death in extreme cases.

Should there be any temporary academic accommodation made for student-athletes who have suffered a concussion?

- Most students will only need help through informal, academic adjustments as they recover from a concussion.
- Students may need to take rest breaks, spend fewer hours at school, be given extra time to complete assignments, as well as being offered other instructional strategies and classroom accommodations
- Contact the school nurse if symptoms persist to discuss whether additional accommodations are



NJSIAA STEROID TESTING POLICY

In accordance with Executive Order 72, issued by the Governor of the State of New Jersey, Richard J. Codey, on December 20, 2005, the NJSIAA will test a random selection of student athletes, who have qualified, as individuals or as members of a team, for state championship competition.

- 1. <u>List of banned substances</u>: A list of banned substances shall be prepared annually by the Medical Advisory Committee and approved by the Executive Committee.
- Consent form: Before participating in interscholastic sports, the student-athlete and the student-athlete's
 parent or guardian shall consent, in writing, to random testing in accordance with this policy. Failure to sign the
 consent form renders the student-athlete ineligible.
- 3. <u>Selection of athletes to be tested</u>: Tested athletes will be selected randomly from all of those athletes participating in championship competition. **Testing may occur at any state championship site or at the school whose athletes have qualified for championship competition.**
- 4. <u>Administration of tests</u>: Tests shall be administered by a certified laboratory, selected by the Executive Director, and approved by the Executive Committee.
- 5. <u>Testing methodology</u>: The methodology for taking and handling samples shall be in accordance with current legal standards.
- 6. <u>Sufficiency of results</u>: No test shall be considered a positive result unless the approved laboratory reports a positive result, and the NJSIAA's medical review officer confirms that there was no medical reason for the positive result.
- 7. <u>Appeal process</u>: If the certified laboratory reports that a student-athlete's sample has tested positive, and the medical review officer confirms that there is no medical reason for a positive result, a penalty shall be imposed unless the student-athlete proves, by a preponderance of the evidence, that he or she bears no fault or negligence for the violation. Appeals shall be heard by a NJSIAA committee consisting of two members of the Executive Committee, the Executive Director/designee, a trainer and a physician. Appeal of a decision of the Committee shall be to the Commissioner of Education, for public school athletes, and to the superior court, for non-public athletes. Hearings shall be held in accordance with NJSIAA By-Laws, Article XIII, "Hearing Procedure."
- 8. <u>Penalties:</u> Any person who tests positively in an NJSIAA administered test, or any person who refuses to provide a testing sample, or any person who reports his or her own violation, shall immediately forfeit his or her eligibility to participate in NJSIAA competition for a period of one year from the date the sample was collected. Any such person shall also forfeit any individual honor earned while in violation. No person who tests positive, refuses to provide a test sample, or who reports his or her own violation shall resume eligibility until he or she has undergone counseling and produced a negative test result.
- 9. <u>Confidentiality</u>: Results of all tests shall be considered confidential and shall only be disclosed to the individual, his or her parents and his or her school.
- 10. <u>Compilation of results</u>: The Executive Committee shall annually compile and report the results of the testing program.
- 11. Yearly renewal of the steroid policy: The Executive Committee shall annually determine whether this policy shall be renewed or discontinued.

Website Resources

- http://tinyurl.com/m2gjmvq Sudden Death in Athletes
- Hypertrophic Cardiomyopathy Association www.4hcm.org
- American Heart Association www.heart.org

Collaborating Agencies:

American Academy of Pediatrics

New Jersey Chapter

3836 Quakerbridge Road, Suite 108 Hamilton, NJ 08619 (p) 609-842-0014 (f) 609-842-0015



American Heart Association

www.aapnj.org

I Union Street, Suite 301 Robbinsville, NJ, 08691 (p) 609-208-0020 www.heart.org



New Jersey Department of Education

www.state.nj.us/education/ Trenton, NJ 08625-0500 (p) 609-292-5935 PO Box 500



New Jersey Department of Health

Frenton, NJ 08625-0360 (p) 609-292-7837 P. O. Box 360

www.state.nj.us/health



Lead Author: American Academy of Pediatrics, New Jersey Chapter

Written by: Initial draft by Sushma Raman Hebbar, MD & Stephen G. Rice, MD PhD

NJ Academy of Family Practice, Pediatric Cardiologists, Additional Reviewers: NJ Department of Education, American Heart Association/New Jersey Chapter, NJ Department of Health and Senior Services, New Jersey State School Nurses

Revised 2014: Nancy Curry, EdM;

Stephen G. Rice, MD; Jeffrey Rosenberg, MD, Lakota Kruse, MD, MPH; Susan Martz, EdM; -ouis Teichholz, MD; Perry Weinstock, MD Christene DeWitt-Parker, MSN, CSN, RN;

ATHLETES CARDIAC SUDDEN **VOUNC**

Sudden Cardiac Death The Basic Facts on in Young Athletes






American Academy of Pediatrics DEDICATED TO THE HEALTH OF ALL CHILDREN"







udden death in young athletes between the ages of 10 done to prevent this kind of What, if anything, can be and 19 is very rare. tragedy?

Res loss hea 엉 Ven 00 ģ anc dn

> What is sudden cardiac death in the young athlete?

ultimately dies unless normal heart rhythm time) during or immediately after exercise heart function, usually (about 60% of the pumping adequately, the athlete quickly result of an unexpected failure of proper is restored using an automated external without trauma. Since the heart stops collapses, loses consciousness, and Sudden cardiac death is the defibrillator (AED).

The an (hi alsc Wit ШП pro ger dev

How common is sudden death in young athletes?

The <u>ō</u>

> Sudden cardiac death in young athletes is The chance of sudden death occurring to any individual high school athlete is reported in the United States per year. very rare. About 100 such deaths are about one in 200,000 per year.

other sports; and in African-Americans than common: in males than in females; in football and basketball than in in other races and ethnic groups. Sudden cardiac death is more

Other diseases of the heart that can lead to sudden death in young people include:

- Myocarditis (my-oh-car-DIE-tis), an acute inflammation of the heart muscle (usually due to a virus).
- Dilated cardiomyopathy, an enlargement of the heart for unknown reasons.
- Long QT syndrome and other electrical abnormalities of the heart which cause abnormal fast heart rhythms that can also run in families.
- Marfan syndrome, an inherited disorder that affects heart valves, walls of major arteries, eyes and the skeleton. It is generally seen in unusually tall athletes, especially if being tall is not common in other family members.

Are there warning signs to watch for?

In more than a third of these sudden cardiac deaths, there were warning signs that were not reported or taken seriously. Warning signs are:

- Fainting, a seizure or convulsions during physical activity;
- Fainting or a seizure from emotional excitement, emotional distress or being startled;
- Dizziness or lightheadedness, especially during exertion;
- Chest pains, at rest or during exertion;
- Palpitations awareness of the heart beating unusually (skipping, irregular or extra beats) during athletics or during cool down periods after athletic participation;
- Fatigue or tiring more quickly than peers; or
- Being unable to keep up with friends due to shortness of breath (labored breathing).

SUDDEN CARDIAC DEATH IN YOUNG ATHLETES What are the current recommendations PPE. However, these procedures

for screening young athletes?

nor

Thi: rev be ath pro can

New Jersey requires all school athletes to be examined by their primary care physician ("medical home") or school physician at least once per year. The New Jersey Department of Education requires use of the specific Preparticipation Physical Examination Form (PPE).

This process begins with the parents and student-athletes answering questions about symptoms during exercise (such as chest pain, dizziness, fainting, palpitations or shortness of breath); and questions about family health history.

The primary healthcare provider needs to know if any family member died suddenly during physical activity or during a seizure. They also need to know if anyone in the family under the age of 50 had an unexplained sudden death such as drowning or car accidents. This information must be provided annually for each exam because it is so essential to identify those at risk for sudden cardiac death.

The required physical exam includes measurement of blood pressure and a careful listening examination of the heart, especially for murmurs and rhythm abnormalities. If there are no warning signs reported on the health history and no abnormalities discovered on exam, no further evaluation or testing is recommended.

Are there options privately available to screen for cardiac conditions?

Technology-based screening programs including a 12-lead electrocardiogram (ECG) and echocardiogram (ECHO) are noninvasive and painless options parents may consider in addition to the required

expensive and are not currently advised by the American Academy of Pediatrics and the American College of Cardiology unless the PPE reveals an indication for these tests. In addition to the expense, other limitations of technology-based tests include the possibility of "false positives" which leads to unnecessary stress for the student and parent or guardian as well as unnecessary restriction from athletic participation.

The United States Department of Health and Human Services offers risk assessment options under the Surgeon General's Family History Initiative available at http://www.hhs.gov/familyhistory/index.html

extrest rest fibrares and but the NJ. "Jar her follower f

When should a student athlete see a heart specialist?

If the primary healthcare provider or school physician has concerns, a referral to a child heart specialist, a pediatric cardiologist, is recommended. This specialist will perform a more thorough evaluation, including an electrocardiogram (ECG), which is a graph of the electrical activity of the heart. An echocardiogram, which is an ultrasound test to allow for direct visualization of the heart structure, will likely also be done. The specialist may also order a treadmill exercise test and a monitor to enable a longer recording of the heart rhythm. None of the testing is invasive or uncomfortable.

Can sudden cardiac death be prevented just through proper screening?

A proper evaluation should find most, but not all, conditions that would cause sudden death in the athlete. This is because some diseases are difficult to uncover and may only develop later in life. Others can develop following a



Keeping Student-Athletes Safe

School athletics can serve an integral role in students' development. In addition to providing healthy forms of exercise, school athletics foster friendships and camaraderie, promote sportsmanship and fair play, and instill the value of competition.

Unfortunately, sports activities may also lead to injury and, in rare cases, result in pain that is severe or long-lasting enough to require a prescription opioid painkiller. It is important to understand that overdoses from opioids are on the rise and are killing Americans of all ages and backgrounds. Families and communities across the country are coping with the health, emotional and economic effects of this epidemic.²

This educational fact sheet, created by the New Jersey Department of Education as required by state law (*N.J.S.A.* 18A:40-41.10), provides information concerning the use and misuse of opioid drugs in the event that a health care provider prescribes a student-athlete or cheerleader an opioid for a sports-related injury. Student-athletes and cheerleaders participating in an interscholastic sports program (and their parent or guardian, if the student is under age 18) must provide their school district written acknowledgment of their receipt of this fact sheet.

How Do Athletes Obtain Opioids?

In some cases, student-athletes are prescribed these medications. According to research, about a third of young people studied obtained pills from their own previous prescriptions (i.e., an unfinished prescription used outside of a physician's supervision), and 83 percent of adolescents had unsupervised access to their prescription medications.³ It is important for parents to understand the possible hazard of having unsecured prescription medications in their households. Parents should also understand the importance of proper storage and disposal of medications, even if they believe their child would not engage in non-medical use or diversion of prescription medications.

What Are Signs of Opioid Use?

According to the National Council on Alcoholism and Drug Dependence, 12 percent of male athletes and 8 percent of female athletes had used prescription opioids in the 12-month period studied.³ In the early stages of abuse, the athlete may exhibit unprovoked nausea and/or vomiting. However, as he or she develops a tolerance to the drug, those signs will diminish. Constipation is not uncommon, but may not be reported. One of the most significant indications of a possible opioid addiction is an athlete's decrease in academic or athletic performance, or a lack of interest in his or her sport. If these warning signs are noticed, best practices call for the student to be referred to the appropriate professional for screening,⁴ such as provided through an evidence-based practice to identify problematic use, abuse and dependence on illicit drugs (e.g., Screening, Brief Intervention, and Referral to Treatment (SBIRT)) offered through the New Jersey Department of Health.


What Are Some Ways Opioid Use and Misuse Can Be Prevented?

According to the New Jersey State Interscholastic Athletic Association (NJSIAA) Sports Medical Advisory Committee chair, John P. Kripsak, D.O., "Studies indicate that about 80 percent of heroin users started out by abusing narcotic painkillers."

The Sports Medical Advisory Committee, which includes representatives of NJSIAA member schools as well as experts in the field of healthcare and medicine, recommends the following:

- The pain from most sports-related injuries can be managed with non-narcotic medications such as acetaminophen, non-steroidal anti-inflammatory medications like ibuprofen, naproxen or aspirin. Read the label carefully and always take the recommended dose, or follow your doctor's instructions. More is not necessarily better when taking an over-the-counter (OTC) pain medication, and it can lead to dangerous side effects.
- Ice therapy can be utilized appropriately as an anesthetic.
- Always discuss with your physician exactly what is being prescribed for pain and request to avoid narcotics.
- In extreme cases, such as severe trauma or post-surgical pain, opioid pain medication should not be prescribed for more than five days at a time;
- Parents or guardians should always control the dispensing of pain medications and keep them in a safe, non-accessible location; and
- Unused medications should be disposed of immediately upon cessation of use. Ask your pharmacist about drop-off locations or home disposal kits like Deterra or Medsaway.

According to NJSIAA Sports
Medical Advisory Committee chair,
John P. Kripsak, D.O., "Studies
indicate that about 80 percent of
heroin users started out by abusing
narcotic painkillers."




NJ Health

STATE OF NEW JERSEY DEPARTMENT OF HEALTH

NISIAA SPORTS MEDICAL ADVISORY COMMITTEE

Karan Chauhan Parsippany Hills High School, Permanent Student Representative New Jersey State Board of Education



Even With Proper Training and Prevention, **Sports Injuries May Occur**

There are two kinds of sports injuries. Acute injuries happen suddenly, such as a sprained ankle or strained back. Chronic injuries may happen after someone plays a sport or exercises over a long period of time, even when applying overuse-preventative techniques.5

Athletes should be encouraged to speak up about injuries, coaches should be supported in injury-prevention decisions, and parents and young athletes are encouraged to become better educated about sports safety.6

What Are Some Ways to Reduce the Risk of Injury?

Half of all sports medicine injuries in children and teens are from overuse. An overuse injury is damage to a bone, muscle, ligament, or tendon caused by repetitive stress without allowing time for the body to heal. Children and teens are at increased risk for overuse injuries because growing bones are less resilient to stress. Also, young athletes may not know that certain symptoms are signs of overuse.

The best way to deal with sports injuries is to keep them from happening in the first place. Here are some recommendations to consider:



PREPARE Obtain the preparticipation physical evaluation prior to participation on a school-sponsored interscholastic or intramural athletic team or squad.



CONDITIONING Maintain a good fitness level during the season and offseason. Also important are proper warm-up and cooldown exercises.



PLAY SMART Try a variety of sports and consider specializing in one sport before late adolescence to help avoid overuse injuries.



ADEQUATE HYDRATION Keep the body hydrated to help the heart more easily pump blood to muscles, which helps muscles work efficiently.



TRAINING Increase weekly training time, mileage or repetitions no more than 10 percent per week. For example, if running 10 miles one week, increase to 11 miles the following week. Athletes should also cross-train and perform sport-specific drills in different ways, such as running in a swimming pool instead of only running on the road.



REST UP Take at least one day off per week from organized activity to recover physically and mentally. Athletes should take a combined three months off per year from a specific sport (may be divided throughout the year in one-month increments). Athletes may remain physically active during rest periods through alternative low-stress activities such as stretching, yoga or walking.



PROPER EQUIPMENT Wear appropriate and properly fitted protective equipment such as pads (neck, shoulder, elbow, chest, knee, and shin), helmets, mouthpieces, face quards, protective cups, and eyewear. Do not assume that protective gear will prevent all injuries while performing more dangerous or risky activities.

Resources for Parents and Students on Preventing Substance Misuse and Abuse

The following list provides some examples of resources:

National Council on Alcoholism and Drug Dependence – NJ promotes addiction treatment and recovery.

New Jersey Department of Health, Division of Mental Health and Addiction Services is committed to providing consumers and families with a wellness and recovery-oriented model of care.

New Jersey Prevention Network includes a parent's quiz on the effects of opioids.

Operation Prevention Parent Toolkit is designed to help parents learn more about the opioid epidemic, recognize warning signs, and open lines of communication with their children and those in the community.

Parent to Parent NJ is a grassroots coalition for families and children struggling with alcohol and drug addiction.

Partnership for a Drug Free New Jersey is New Jersey's anti-drug alliance created to localize and strengthen drug-prevention media efforts to prevent unlawful drug use, especially among young people.

The Science of Addiction: The Stories of Teens shares common misconceptions about opioids through the voices of teens.

Youth IMPACTing NJ is made up of youth representatives from coalitions across the state of New Jersey who have been impacting their communities and peers by spreading the word about the dangers of underage drinking, marijuana use, and other substance misuse.

- References ¹ Massachusetts Technical Assistance Partnership for Prevention
 - ² Centers for Disease Control and Prevention
 - ³ New Jersey State Interscholastic Athletic
- Association (NJSIAA) Sports Medical Advisory Committee (SMAC)
- 4 Athletic Management, David Csillan, athletic trainer, Ewing High School, NJSIAA SMAC
- 5 National Institute of Arthritis and Musculoskeletal and Skin Diseases
- ⁶ USA TODAY
- 7 American Academy of Pediatrics

An online version of this fact sheet is available on the New Jersey Department of Education's Alcohol, Tobacco, and Other Drug Use webpage. Updated Jan. 30, 2018.

SPORTS-RELATED EYE INJURIES:

AN EDUCATIONAL FACT SHEET FOR PARENTS





Participating in sports and recreational activities is an important part of a healthy, physically active lifestyle for children. Unfortunately, injuries can, and do, occur. Children are at particular risk for sustaining a sports-related eye injury and most of these injuries can be prevented. Every year, more than 30,000 children sustain serious sports-related eye injuries. Every 13 minutes, an emergency room in the United States treats a sports-related eye injury. According to the National Eye Institute, the sports with the highest rate of eye injuries are: baseball/softball, ice hockey, racquet sports, and basketball, followed by fencing, lacrosse, paintball and boxing.

Thankfully, there are steps that parents can take to ensure their children's safety on the field, the court, or wherever they play or participate in sports and recreational activities.

Prevention of Sports-Related Eye Injuries

Approximately 90% of sports-related eye injuries can be prevented with simple precautions, such as using protective eyewear.² Each sport has a certain type of recommended protective eyewear, as determined by the American Society for Testing and Materials (ASTM). Protective eyewear should sit comfortably on the face. Poorly fitted equipment may be uncomfortable, and may not offer the best eye protection. Protective eyewear for sports includes, among other things, safety goggles and eye guards, and it should be made of polycarbonate lenses, a strong, shatterproof plastic. Polycarbonate lenses are much stronger than regular lenses.³

Health care providers (HCP), including family physicians, ophthalmologists, optometrists, and others, play a critical role in advising students, parents and guardians about the proper use of protective eyewear. To find out what kind of eye protection is recommended, and permitted for your child's sport, visit the National Eye Institute at http://www.nei.nih.gov/sports/findingprotection.asp. Prevent Blindness America also offers tips for choosing and buying protective eyewear at http://www.preventblindness.org/tips-buying-sports-eye-protectors, and http://www.preventblindness.org/ recommended-sports-eye-protectors.

It is recommended that all children participating in school sports or recreational sports wear protective eyewear. Parents and coaches need to make sure young athletes protect their eyes, and properly gear up for the game. Protective eyewear should be part of any uniform to help reduce the occurrence of sports-related eye injuries. Since many youth teams do not require eye protection, parents may need to ensure that their children wear safety glasses or goggles whenever they play sports. Parents can set a good example by wearing protective eyewear when they play sports.

¹ National Eye Institute, National Eye Health Education Program, Sports-Related Eye Injuries: What You Need to Know and Tips for Prevention, www.nei.nih.gov/sports/pdf/sportsrelatedeyeInjuries.pdf, December 26, 2013.

² Rodriguez, Jorge O., D.O., and Lavina, Adrian M., M.D., Prevention and Treatment of Common Eye Injuries in Sports, http://www.aafp.org/afp/2003/0401/p1481.html, September 4, 2014; National Eye Health Education Program, Sports-Related Eye Injuries: What You Need to Know and Tips for Prevention, www.nei.nih.gov/sports/pdf/sportsrelatedeyeInjuries.pdf, December 26, 2013.

³ Bedinghaus, Troy, O.D., Sports Eye Injuries, http://vision.about.com/od/emergencyeyecare/a/Sports_Injuries.htm, December 27, 2013.

Most Common
Types of Eye
Injuries

The most common types of eye injuries that can result from sports injuries are blunt injuries, corneal abrasions and penetrating injuries.

- ◆ Blunt injuries: Blunt injuries occur when the eye is suddenly compressed by impact from an object. Blunt injuries, often caused by tennis balls, racquets, fists or elbows, sometimes cause a black eye or hyphema (bleeding in front of the eye). More serious blunt injuries often break bones near the eye, and may sometimes seriously damage important eye structures and/or lead to vision loss.
- ◆ Corneal abrasions: Corneal abrasions are painful scrapes on the outside of the eye, or the cornea. Most corneal abrasions eventually heal on their

own, but a doctor can best assess the extent of the abrasion, and may prescribe medication to help control the pain. The most common cause of a sports-related corneal abrasion is being poked in the eye by a finger.

- ◆ Penetrating injuries: Penetrating injuries are caused by a foreign object piercing the eye. Penetrating injuries are very serious, and often result in severe damage to the eye. These injuries often occur when eyeglasses break while they are being worn. Penetrating injuries must be treated quickly in order to preserve vision.⁴
- Pain when looking up and/or down, or difficulty seeing;
- Tenderness;
- Sunken eye;
- Double vision;
- Severe eyelid and facial swelling;
- Difficulty tracking;

Signs or Symptoms of an Eye Injury



- The eye has an unusual pupil size or shape;
- · Blood in the clear part of the eye;
- Numbness of the upper cheek and gum; and/or
- Severe redness around the white part of the eye.

What to do if a Sports-Related Eye Injury Occurs

If a child sustains an eye injury, it is recommended that he/she receive immediate treatment from a licensed HCP (e.g., eye doctor) to reduce the risk of serious damage, including blindness. It is also recommended that the child, along with his/her parent or guardian, seek guidance from the HCP regarding the appropriate amount of time to wait before returning to sports competition or practice after sustaining an eye injury. The school nurse and the child's teachers should also be notified when a child sustains an eye injury. A parent or guardian should also provide the school nurse with a physician's note detailing the nature of the eye injury, any diagnosis, medical orders for

the return to school, as well as any prescription(s) and/or treatment(s) necessary to promote healing, and the safe resumption of normal activities, including sports and recreational activities.

According to the American Family Physician Journal, there are several guidelines that should be followed when students return to play after sustaining an eye injury. For

Return to Play and Sports

example, students who have sustained significant ocular injury should receive a full examination and clearance by an ophthalmologist or optometrist. In addition, students should not return to play until the period of time recommended by their HCP has elapsed. For more minor eye injuries, the athletic trainer may determine that

it is safe for a student to resume play based on the nature of the injury, and how the student feels. No matter what degree of eye injury is sustained, it is recommended that students wear protective eyewear when returning to play and immediately report any concerns with their vision to their coach and/or the athletic trainer.

Additional information on eye safety can be found at http://isee.nei.nih.gov and http://www.nei.nih.gov/sports.