HEALTH & PHYSICAL EDUCATION DEPARTMENT



Mr. Nephtaly Cardona
Director of Athletics, Health and Physical Education (9-12)
Perth Amboy Public Schools
300 Eagle Avenue
732-376-6030 Ext. 23-408

nephcardona@paps.net

In preparation for the reopening of athletics, the following state approved athletic physical forms can now be **downloaded online through the <u>www.paps.net</u> website**; accessed in the **Athletics/Health Services tab**. Reminder that all physical forms must be completed and submitted no later than Wednesday **July 22nd 2020**. Fall Season begins on Monday **July 27th 2020**.

Sports Physical Forms Distribution

(You may pick up a blank physical packet at these locations below)

Location	Dates	Time
Rose M. Lopez School	Monday-Wednesday-Friday	8:30am-11:30pm
Robert N. Wilentz School	Monday-Wednesday-Friday	8:30am-11:30pm
PAHS Main Campus	Monday-Wednesday-Friday	8:30am-11:30pm

The following FREE Sports Physical Dates listed are for all student-athletes who wish to participate during the <u>Fall Season Only</u>. Student must bring the following forms completed and signed by their parent/guardian in order to receive the FREE Physical.

- History Form
- o COVID- 19 Questionnaire
- o Sudden Cardiac Death Sign Off
- o Opioid Drugs Sign Off
- NJSIAA Steroid Consent
- Concussion Sign Off

FREE Fall Sports Physical Dates

(First Come, First Serve)

Location	Dates	Times
Dr. Herbert N. Richardson School	Saturday, July 11, 2020	9am-12pm
	Wednesday, July 15, 2020	10am-1pm
(318 Stockton Street	Saturday, July 18, 2020	9am-12pm
Perth Amboy, NJ 08861)	Wednesday July 22, 2020	10am-1pm

Sports Physical Forms Collection/Drop-Off

(If you are receiving a physical with your private physician, please drop off your completed physical packet at the following location during the dates and times listed)

Location	Dates	Times
Dr. Herbert N. Richardson School	Saturday, July 11, 2020	9am-12pm
	Wednesday, July 15, 2020	10am-1pm
(318 Stockton Street	Saturday, July 18, 2020	9am-12pm
Perth Amboy, NJ 08861)	Wednesday July 22, 2020	10am-1pm

ALL FORMS ARE DUE BY WEDNESDAY JULY 22nd @ Dr. Herbert N. Richardson School

ATTENTION PARENT/GUARDIAN: The preparticipation physical examination (page 3) must be completed by a health care provider who has completed the Student-Athlete Cardiac Assessment Professional Development Module.

M PREPARTICIPATION PHYSICAL EVALUATION HISTORY FORM

۵			Date of birth		
Ann Grado Schoo	1		Sport(s)		
dicines and Allergies: Please list all of the prescription and over-tr	1e-cou	nter me	dictnes and supplements (herbal and nutritional) that you are currently t	alung	
		-			
	•	-10 Da			
you have any atlergies? Yes No if yes, pleaso identifyed Pollens	ry spe	CINC SUE)	rgy delew. □ Food □ Stinging insects		
					•
ain "Yes" answers below, Circle questions you don't know the answ	Yes	Eto I	MEDICAL QUESTIONS	Yes	Mo
HERAL QUESTIONS Has a doctor ever denied or restricted your participation in sports for	163		26. On you cough, wheeve, or have difficulty breathing during or		
SUA LESTOUS.			after exercise?		
Do you have any enpoine medical conditions? If so, please identify			27. Have you ever used an inhater or taken asihma medicine?		
below: Asthma Anemia Diabetes Infections		- 1	St. is there anyone in your facility who has asthme? Wate you born without or are you missing a kidney, an eye, a testide		\vdash
Uner:			(majes), your spices, or any other organ?		
Have you ever about the ratin or an incoherent			30. Do you have grain pain or a painful bulge or hernia in the grain area?		
ART HEALTH QUESTIONS ABOUT YOU	Yes	No	31. Have you had infectious monanucleosis (mone) within the last month?		L
Have you ever passed out or nearly passed out DURING or			32. Do you have any rashes, pressure seres, or other skin problems?		<u> </u>
AFTER exercise?			33. Have you had a herpes or MRSA skin infection?		
Hove you ever had discomfort, pain, tightness, or pressure in your			34. Have you ever had a head injury or concussion?		├
chast during exercise? Does your heart ever race or skip beats (fregular beats) during exercise?			35. Have you over had a hit or blow to the head that caused confusion, protonged hazdache, or memory problems?		
. Has a doctor ever told you that you have any heart problems? If so,			36. Do you have a history of scizure disorder?		I^-
check all that apply:		1	37. Do you have headaches with exercise?		
☐ High blood pressure ☐ A heart murmur			38. Have you over had numbross, tingling, or weakness in your arms or		Π
High cholestorol A heart infection Kawasaki disease Other:			tees after being hit or falling?		╁
Has a doctor ever ordered a test for your heart? (For oxample, ECG/EKG.			39. Have you ever been unable to move your urms or legs after being hit or falling?		1
echocardiogram)			40. Have you ever became ill white exercising in the heat?		
Do you get Eghtheaded or feel more short of breath than expected			41. Do you get frequent muscle cramps when exercising?		
during exercise? Have you ever had an unexplained selzure?		-	42. Do you or someone in your family have sickle cell trait or disease?		L
. Have you ever has an energisened sectors? Do you get more tired or short of breath more quickly than your lifends		 -	43. Have you had any problems with your eyes or vision?	<u> </u>	上
during exercise?		<u></u>	44. Have you had any eye injuries?		丄
EART HEALTH QUESTIONS ABOUT YOUR FAMILY	Yes	Ro	45. Do you wear glasses or contact lenses?	 	1
. Has any family member or relative died of heart problems or had an			46. Do you wear protective eyewear, such as goggles or a face shield?	├	╀
unexpected or unexplained sudden death before age 50 (including drowning, unexplained car accident, or sudden infant death syndrome)?]	47. Do you warry about your weight?	├	╁╌
A Commence to what to the home transferrable existing the Marian			48. Are you trying to or has anyone recommended that you gain or lase welch?		1
and a second bases of a second contribution of the contribution of			49. Are you on a special diet or do you avoid certain types of foods?	1	
syndrome, short OT syndrome, brugada syndrome, or catecholaminorgic polymorphic ventricular tachycardia?			50. Have you ever had an eating disorder?		1
5. Does enyone in your family have a heart problem, pacemaker, or			51. Do you have any concerns that you would like to discuss with a doctor?		L
implanted delibrilistor?	<u></u> _	┼	FEMALES CHLY	<u> </u>	┸
6. Has anyone in your family had unexplained telaling, unexplained seizuros, or not drowning?			52. Have you ever had a menstrual period?		
SEARCE, OF THAT CHESTICHS	Yes	Ho	53. How old were you when you had your first menstrual period?	┼	
7. Have you ever had an injury to a bone, muscle, Egament, or tenden			54. How many periods have you had in the last 12 menths?		
that caused you to miss a practice or a game?	├	↓ —	Expisin "yes" answers here		
8. Have you ever had any broken or fractured bones or distocated joints?	 -	┼			
Have you over had an injury that required x-rays, MRI, CT scen. Injections, therapy, a brace, a cast, or cruiches?					
ryccions, userapy, a urace, a case, or country. 10. Have you ever had a stress fracture?		1			
to Name were more been told that you have or have you had an x-ray for neck	1	1			
instability or attenteaxial instability? (Down syndrome or owarrism)	↓	4-			
22. Do you regularly use a brace, ortholics, or other assistive device?	 -	┼			
23. Do you have a bone, muscle, or joint injury that bothers you?	 	╁			
24. Do any oi your joints become painful, swellen, feel warm, or look red? 25. Do you have any history of Juvenile arthritis or connective tissue disease?	1-	+			
no and the company of branch admitted of connection there is discussed.	1				

© 2010 American Academy of Family Physicians, American Academy of Podistrics, American College of Sports Medicine, American Med

M PREPARTICIPATION PHYSICAL EVALUATION THE ATHLETE WITH SPECIAL NEEDS: SUPPLEMENTAL HISTORY FORM

Date of Exam						
Name				Date of birth	n	
	Ana	Grada	School	Sport(s)		
26x	. Age	Graus	3CH001	openior		
1. Type of dis	ability					
2. Date of cis	ability					· · · · · · · · · · · · · · · · · · ·
3. Classificati	on (il avallable)					• •
4. Causo of 6	inability (birth, dis	sease, accident/trauma, cihsi)			
	orts you are inter		· · · · · · · · · · · · · · · · · · ·			
					Yes	Ho
6. Do you reg	ularly uso a brac	e, assistive device, or prosthe	stic?			
7. Do you use	any special brac	ce or assistive device for spor	ts?			<u> </u>
8. Do you have	ro any rashes, pri	assure sores, or any other ski	in problems?			
9. Do you have	re a hearing leas	? Do you use a boaring aid?				<u> </u>
10. Do you hav	ra a visual impeli	rment?				
11. Do you use	any special devi	ices for bowel or bladder fund	ction?			
12. Do you han	re burning or disc	comfort when urinating?				
13. Have you l	nad autonomic dy	rsreflexia?				٠.
14. Have you o	ver been dizgno	sed with a heat-related (hype	rihermia) or cold-related (hypothermia) iline	95?		<u> </u>
	ro musclo spastic					
16. Do you han	ro frequent seizu	res that cannot be controlled	by medication?			<u> </u>
Explain "yes" s	nawers here					
•						
						
						• •
EXECUTED HORIZONIA	O II YOU CENE ON	er had any of the following.				
12000					Yes	No
	in the second				Yes .	No
Atlantoaxial In:	elability				Yes	No
Atlantoaxial in: X-ray ovaluation	stability on for atlantoaxia	l instability			Yes	No
Atlantoaxial in: X-ray ovaluation	elability	l instability			Yes .	No
Atlantoaxial in: X-ray ovaluatio Dislocated join Easy bloeding	stability on for allantoaxia ds (more than on	l instability			Yes	No
Atlantoaxial in: X-ray ovaluation Distocated join Easy blooding Enlarged splco	stability on for allantoaxia ds (more than on	l instability			Yes	No
Atlantoaxial in: X-ray ovaluation Dislocated join Easy bloeding Enlarged spice Hepatitis	stability on for atlantosxia its (more than on	I instability			Yes	No
Atlantoaxisi in: X-ray ovaluatio Dislocated join Easy bleeding Entarged spice Hepatitis Ostoopenia or	stability on for allantosxial its (more than on in	I instability			Yes	No
Atlantoaxial in: X-ray ovaluation Uslocated John Easy blaeding Enlarged splce Hepatitis Ostoopenia or Difficulty contr	stability on for attantoaxia its (more than on the osteoporosis relling boxel	I instability			Yes	No
Atlantoaxial in: X-ray ovaluatio Distocated join Easy blooding Entarged spice Hepatitis Oatoopenia or Officulty contr	stability on for allantoaxia is (more than on on osteoporosis relling bowel relling bladder	l instability e)			Yes	No
Atlantoaxial in: X-ray ovaluatio Distocated join Easy blaeding Entarged spice Hepatitis Ostoopenia or Otificuity costs Numbress or	stability on for atlantosxia is (more than on osteoporosis relling bowel relling bladder lingling in arms o	i instability e) or hands			Yes	No
Atlantoaxial in: X-ray ovaluatio Distocated join Easy bloeding Entarged spice Hepatitis Ostoopenia or Otificuity costs Numbness or Numbness or	stability on for attantoaxia is (more than on osteoporosis relling bowel relling bladder lingling in arms o tingling in legs or	i instability e) or hands			Yes	No
Atlantoaxial in: X-ray ovaluatio Distocated join Easy bloeding Entarged spice Hepatitis Ostoopenia or Otificuity costs Numbness or Numbness or Weskness in a	stability on for attentoaxia is (more than on osteoporosis rating bowel rating bladder lingling in arms o tingling in legs or orms or bands	i instability e) or hands			Yes	No
Atlantoaxial in: X-ray ovaluatio Distocated join Easy blaeding Entarged spice Hepatitis Ostoopenia or Otificuity control Numbness or Numbness or Westness in a Westness in it	stability in for attantoaxia is (more than on in osteoporosis relling bowel relling bladder lingling in arms o tingling in legs or urms or bands egs or feet	i instability e) or hands r feet			Yes	No
Atlantoaxial in: X-ray ovaluatio Distocated join Easy blaeding Entarged spice Hepatitis Ostoopenia or Otificuity costs Numbness or Numbness or Westness in a Weatness in it Recent change	stability on for attentoaxia is (more than on osteoporosis relling bowel relling bladder lingling in arms o tingling in legs or orms or bands egs or feet a in coordination	I instability o) or hands or feet			Yes	No
Atlantoaxial in: X-ray ovaluatio Distocated join Easy blaeding Entarged splce Hepatitis Ostoopenia or Otificuity costs Numbness or Numbness or Westness in a Weatness in it Recent change	stability in for attantoaxia is (more than on in osteoporosis relling bowel relling bladder lingling in arms o tingling in legs or urms or bands egs or feet	I instability o) or hands or feet			Yes	No
Atlantoaxial in: X-ray ovaluatio Distocated join Easy blaeding Entarged spice Hepatitis Ostoopenia or Otificuity costs Numbness or Numbness or Westness in a Westness in b Recent change Spina bifido	stability on for attentoaxia is (more than on osteoporosis relling bowel relling bladder lingling in arms o tingling in legs or orms or bands egs or feet a in coordination	I instability o) or hands or feet			Yes	No
Attentoaxist in: X-ray ovatuation Essy blooding Enlarged spice Hepatities Ostopenia or Difficulty control Numbness or in Westness in a Weatness in a Weatness in a Recent change Rocent change Spina blifidu Letex allergy	stability on for allantosxial is (more than on in estemporosis relling bowel relling bladder rellinging in logs or ingring in l	I instability o) or hands or feet			Yes	No
Atlantoaxial in: X-ray ovaluatio Distocated join Easy blaeding Entarged spice Hepatitis Ostoopenia or Otificuity costs Numbness or Numbness or Westness in a Westness in b Recent change Spina bifido	stability on for allantosxial is (more than on in estemporosis relling bowel relling bladder rellinging in logs or ingring in l	I instability o) or hands or feet			Yes	No
Attentoaxist in: X-ray ovatuation Essy blooding Enlarged spice Hepatities Ostopenia or Difficulty control Numbness or in Westness in a Weatness in a Weatness in a Recent change Rocent change Spina blifidu Letex allergy	stability on for allantosxial is (more than on in estemporosis relling bowel relling bladder rellinging in logs or ingring in l	I instability o) or hands or feet			Yes	No
Attentoaxist in: X-ray ovatuation Essy blooding Enlarged spice Hepatities Ostopenia or Difficulty control Numbness or in Westness in a Weatness in a Weatness in a Recent change Rocent change Spina blifidu Letex allergy	stability on for allantosxial is (more than on in estemporosis relling bowel relling bladder rellinging in logs or ingring in l	I instability o) or hands or feet			Yes	
Attentoaxist in: X-ray ovatuation X-ray ovatuation Essy blooding Enlarged spice Hepatities Ostopenia or Difficulty control Numbness or in Numbness or in Weskness in a Weskness in a Weskness in a Recent change Rocent change Spina blifidu Letex allergy	stability on for allantosxial is (more than on in estemporosis relling bowel relling bladder rellinging in logs or ingring in l	I instability o) or hands or feet			Yes	
Attentoaxist in: X-ray ovatuation X-ray ovatuation Essy blooding Enlarged spice Hepatities Ostopenia or Difficulty control Numbness or in Numbness or in Weskness in a Weskness in a Weskness in a Recent change Rocent change Spina blifidu Letex allergy	stability on for allantosxial is (more than on in estemporosis relling bowel relling bladder rellinging in logs or ingring in l	I instability o) or hands or feet			Yes	
Attentoaxist in: X-ray ovatuation X-ray ovatuation Essy blooding Enlarged spice Hepatities Ostopenia or Difficulty control Numbness or in Numbness or in Weskness in a Weskness in a Weskness in a Recent change Rocent change Spina blifidu Letex allergy	stability on for allantosxial is (more than on in estemporosis relling bowel relling bladder rellinging in logs or ingring in l	I instability o) or hands or feet			Yes	
Attentoaxist in: X-ray ovatuation X-ray ovatuation Essy blooding Enlarged spice Hepatities Ostopenia or Difficulty control Numbness or in Numbness or in Weskness in a Weskness in a Weskness in a Recent change Rocent change Spina blifidu Letex allergy	stability on for allantosxial is (more than on in estemporosis relling bowel relling bladder rellinging in logs or ingring in l	I instability o) or hands or feet			Yes	
Atlantoaxial in: X-ray ovaluatic Unitocated join Easy blaeding Enlarged spice Hepatitis Ostoopenia or Difficulty contr Difficulty contr Numbness or Numbness or Weakness in it Recent change Recent change Spina bifidu Latex allergy Explain "yes"	stability In for allantoaxia is (more than on in esteaporoxis reling bowel reling bladder lingling in arms or tingling in legs or arms or bands egs or feet a in coordination o in ability to wall acswers here	i instability e) w hands r feet		a and correct.	Yes	
Atlantoaxial in: X-ray ovaluatic Unitocated join Easy bloeding Entarged spice Hepatitis Ostoopenia or Otificuity costs Numbness or Numbness or Weakness in a Weakness in it Recent change Recent change Spina bifidu Latex allergy Explain "yes"	stability In for allantoaxia is (more than on in esteaporoxis reling bowel reling bladder lingling in arms or tingling in legs or arms or bands egs or feet a in coordination o in ability to wall acswers here	i instability e) w hands r feet		a and correct.	Yes	

NOTE: The preparticiaption physical examination must be conducted by a health care provider who 1) is a licensed physician, advanced practice nurse, or physician assistant; and 2) completed the Student-Athlete Cardiac Assessment Professional Development Module.

_____ Date of birth _

PREPARTICIPATION PHYSICAL EVALUATION PHYSICAL EXAMINATION FORM

Name

PHYSICIAN REMINDERS

Do you feel stressed out or under a lot of pressure? Do you ever feel sod, hopeless, depressed, or analous? So you feel sode at your home or residence?			
* Have you ever tried eigeneties, chewing tobacco, snuff, or dip? * Buring the past 30 days, did you use chewing tobacco, snuff, or dip?			
 Do you drink alcohol or use any other drugs? 			
* Have you exact taken anyboile steroids or used any other performance supplement?	navlannonna9		
 Have you ever taken any supplements to help you gain or lose weight or improve your Do you weer a seat bell, uso a helmet, and use condoms? 	bester instances		
Consider roviewing questions on cardiovascular symptems (questions 5–14).			
ROITANIMAX			
alght Weight D Male	☐ Female		
P / (/) Pulse Vision		L 20/	Corrected CI Y CI N
EDICAL	NORMAL		ABKORMAL FINDINGS
ppezrance Marian stigmata (hyphosoziosis, high-arched polate, poctus oxcavatum, arachnedactyly, arm spon > hajahi, hyperkodly, myopia, MVP, cordo insufficiency)			
yes/ears/nose/hrost Pupils equal			
Hearing	ļ		
mph nodes	 		
Aturmurs (auscutation standing, supine, +/- Valsatva) Location of point of maximal imputse (PKI)			
ulses Simultaneous femoral and radial pulses	<u> </u>		
2003 2000012022018 (Brutol 8) firit (Batris) Entrace	1		
bdomen			
enitourinary (maies only)*			
kin		İ	
HSV, lesions suggestive of MRSA, lines corports eurologic*			
RUSCUILOS KELETAL			
lock			
ack			
haufder/arm	ļ		
lt.pre/forearm	╂		
Vrist/hand/lingers			
lip/thigh (nee			
ebina/aa			
outroes			
runcilonál • Duck-welk, single leg hop			
onsider ECC, echocardiogram, and reternal to cardiology for abnormal cardiac history or exam. consider GD ream B in private action. Having birling privates its recommended. consider cognitive evaluation or bosciles resuspsychiatric testing if a history of significant concursion.			
Cleared for all sports without restriction			
Cleared for all sports without restriction with recommendations for further evaluation or treats	пети юг		
Not cleared			
Pending further evaluation			
☐ For any sports			
□ For certain sports			
Rassa	<u>, , , , , , , , , , , , , , , , , , , </u>		
ecommandations			
nave examined the above-named student and completed the preparticipation physical e utilizipate in the sperife) as culticad above. A copy of the physical exam is on record in m ise after the sthicte has been cleared for participation, a physician may reseted the clear	n ed nes bae esillo v	lade avallable to	(DB 20000) Sf (IIS LEGIBAL OI IIIS BELSIIIS" II COINCI
ise athiete (and parents/guardians).			
tome of obsciring advanced practice purse (APN), physician assistant (PA) (print/type)_			Date of exam
Address			Phone
Signature of physician, APN, PA			
The state of the s			
2010 American Academy of Family Physicians, American Academy of Pediatrics, American CoS	oge of Sports Medicine	American Medica	l Society for Sports Medicino, American Orthopoedic
lociety for Sports titedicine, and American Osteopathic Academy of Sports Medicine. Permission	is granted to reprint fo	r nancommercial, e	ducational purposes with acknowledgment. 9-8-
tetio haw Jersey Department of Education 2014; Pursuant to P.L.2013, c.71			140

M PREPARTICIPATION PHYSICAL EVALUATION CLEARANCE FORM

Name		Sex 🗆 M	O F	Age	Date of birth
☐ Cleared for a	ail sports without restriction				
Cleared for	ed sports wilhout restriction with recommendations for further eva	duation or trea	tment f	or	
Not cleared					
0	Pending further evaluation				•
	For any sports				
	For certain sports				
	Reason				
Recommendation	ons	-			
					<u> </u>
EMERGENC	CY INFORMATION				
					••
					•
-					
Other informati	ng .				
Oftest menusan					
					
HAD APPLICE C	YAND	SCHO	OL PH	YSICIAN:	•
HCP OFFICE S	IAMP	1 —			
		1 1		· -	(Date)
		Ap	prove	d	Not Approved
		Sig	natur	e:	
		ـا ل			
I have exam	ined the above-named student and completed the preparations to practice and participate in the sport(s	participation	ı physi	ical evalu	ation. The athlete does not present apparent of the physical exam is on record in my office
and any he	made available to the cohool at the reguest of the HRIE	nts. if condi	tions a	arise ante	L TUB STUTETS USS DESU CLEATER IN hatricihannii
the physicia	in may rescind the clearance until the problem is resol	ved and the	poten	tial cons	equences are completely explained to the athlet
(and parent	s/guardians).				
Name of phys	sician, advanced practice nurse (APN), physician assistant (P/	A)			Oals
· · · · · · · · · · · · · · · · · · ·	hysician, APN, PA				
•	ardiac Assessment Professional Development Module				
	Signature	=			
O016					

©2010 American Academy of Family Physicians, American Academy of Pediatrics, American College of Sports Medicine, American Electrical Society for Sports Medicine, American Ostropadhic Academy of Sports Medicine. Permission is granted to reprint for noncommercial, educational purposes with acknowledgment.

New Jersey Department of Education 2014; Pursuant to P.L.2013, c.71

NEW JERSEY STATE INTERSCHOLASTIC ATHLETIC ASSOCIATION

1161 Route 130 North, Robbinsville, NJ 08691-1104 Phone 609-259-2776 ~ Fax 609-259-3047

COVID-19 Questionnaire

Name of Student:	Date:	
Parent/Guardian Cell:	Sport:	
COVID-19 Questions:	Please Ci	rcle One
Has your son/daughter been diagnosed with Coronavirus (COVID-19)?	YES	NO
x If diagnosed with Coronavirus (COVID-19), was your son/daughter symptomatic?	YES	NO ·
x If diagnosed with Coronavirus (COVID-19), was your son/daughter hospitalized?	YES	NO
Has any member of the student-athlete's household been diagnosed with Coronavirus (COVID-19)?	YES	NO
Signature of Parent/Guardian:		
To participate in workouts during the summer recess period, the parent/guardia	n <u>must</u> complete th	is form.
Student ID#		٠,
Student Name -		•

State of New Jersey DEPARTMENT OF EDUCATION

Sudden Cardiac Death Pamphlet Sign-Off Sheet

Name of School District:
Name of Local School:
I/We acknowledge that we received and reviewed the Sudden Cardiac Death in Young Athletes pamphlet.
Student Signature:
Parent or Guardian Signature:
Date:

Use and Misuse of Opioid Drugs Fact Sheet

Student-Athlete and Parent/Guardian Sign-Off

In accordance with N.J.S.A. 18A:40-41.10, public school districts, approved private schools for students with disabilities, and nonpublic schools participating in an interscholastic sports program must distribute this Opioid Use and Misuse Educational Fact Sheet to all student-athletes and cheerleaders. In addition, schools and districts must obtain a signed acknowledgement of receipt of the fact sheet from each student-athlete and cheerleader, and for students under age 18, the parent or guardian must also sign.

	Name of School:
	Name of School District (if applicable): Perth Amboy Public Schools
	I/We acknowledge that we received and reviewed the Educational Fact Sheet on the Use and Misuse of Opioid Drugs.
*	Student Signature:
*	Parent/Guardian Signature (also needed if student is under age 18):

Date:

£

NJSIAA

1161 Route 130, P.O. Box 487, Robbinsville, NJ 08691

609-259-2776 609-259-3047-Fax

NJSIAA STEROID TESTING POLICY

CONSENT TO RANDOM TESTING

In Executive Order 72, issued December 20, 2005, Governor Richard Codey directed the New Jersey Department of Education to work in conjunction with the New Jersey State Interscholastic Athletic Association (NJSIAA) to develop and implement a program of random testing for steroids, of teams and individuals qualifying for championship games.

Beginning in the Fall, 2006 sports season, any student-athlete who possesses, distributes, ingests or otherwise uses any of the banned substances on the attached page, without written prescription by a fully-licensed physician, as recognized by the American Medical Association, to treat a medical condition, violates the NJSIAA's sportsmanship rule, and is subject to NJSIAA penalties, including ineligibility from competition.

Athletes may submit supplements and medications to Drug Free Sport AXIS to receive information regarding banned substances or safety issues. Athletes or parents may login to the NJSIAA account at www.dfsaxis.com using the password "njsports".

The NJSIAA will test certain randomly selected individuals and teams that qualify for a state championship tournament or state championship competition for banned substances. The results of all tests shall be considered confidential and shall only be disclosed to the student, his or her parents and his or her school. No student may participate in NJSIAA competition unless the student and the student's parent/guardian consent to random testing.

By signing below, we consent to random testing in accordance with the NJSIAA steroid testing policy. We understand that, if the student or the student's team qualifies for a state championship tournament or state championship competition, the student may be subject to testing for banned substances.

Signature of Student-Athlete	Print Student-Athlete's Name	Date
Signature of Parent/Guardian	Print Parent/Guardian's Name	Date

What Should a Student-Athlete do if they think they have a concussion?

- Don't hide it. Tell your Athletic Trainer, Coach, School Nurse, or Parent/Guardian.
- Report it. Don't return to competition or practice with symptoms of a concussion or head injury. The sooner you report it, the sooner you may return-to-play.
- Take time to recover. If you have a concussion your brain needs time to heal. While your brain is
 healing you are much more likely to sustain a second concussion. Repeat concussions can cause
 permanent brain injury.

What can happen if a student-athlete continues to play with a concussion or returns to play to soon?

- Continuing to play with the signs and symptoms of a concussion leaves the student-athlete vulnerable to second impact syndrome.
- Second impact syndrome is when a student-athlete sustains a second concussion while still having symptoms from a previous concussion or head injury.
- Second impact syndrome can lead to severe impairment and even death in extreme cases.

Should there be any temporary academic accommodations made for Student-Athletes who have suffered a concussion?

- To recover cognitive rest is just as important as physical rest. Reading, texting, testing-even watching
 movies can slow down a student-athletes recovery.
- Stay home from school with minimal mental and social stimulation until all symptoms have resolved.
- Students may need to take rest breaks, spend fewer hours at school, be given extra time to complete
 assignments, as well as being offered other instructional strategies and classroom accommodations.

Student-Athletes who have sustained a concussion should complete a graduated return-to-play before they may resume competition or practice, according to the following protocol:

- Step 1: Completion of a full day of normal cognitive activities (school day, studying for tests, watching
 practice, interacting with peers) without reemergence of any signs or symptoms. If no return of symptoms,
 next day advance.
- Step 2: Light Aerobic exercise, which includes walking, swimming, and stationary cycling, keeping the
 intensity below 70% maximum heart rate. No resistance training. The objective of this step is increased
 heart rate.
- Step 3: Sport-specific exercise including skating, and/or running: no head impact activities. The objective
 of this step is to add movement.
- Step 4: Non contact training drills (e.g. passing drills). Student-athlete may initiate resistance training.

For further information on Sports Polated Concussions and other Head Injuries please visit:

- Step 5: Following medical clearance (consultation between school health care personnel and studentathlete's physician), participation in normal training activities. The objective of this step is to restore confidence and assess functional skills by coaching and medical staff.
- Step 6: Return to play involving normal exertion or game activity.

bianj.org www.atsnj.org	
' Ct. J Athlete's Nome	Data
rint Student-Athlete's Name	Date
	Date
	Print Parent/Guardian's Name

2020-21 NJSIAA Banned Substances

It is the student-athlete's responsibility to check with the appropriate or designated athletics staff before using any substance.

The NJSIAA bans the following drug classes.

- a. Stimulants.
- b. Anabolic agents.
- c. Alcohol and beta blockers.
- d. Diuretics and masking agents.
- e. Narcotics.
- f. Cannabinoids.
- g. Peptide hormones, growth factors, related substances and mimetics.
- h. Hormone and metabolic modulators (anti-estrogens).
- i. Beta-2 agonists.

Note: Any substance chemically/pharmacologically related to all classes listed above and with no current approval by any governmental regulatory health authority for human therapeutic use (e.g., drugs under pre-clinical or clinical development or discontinued, designer drugs, substances approved only for veterinary use) is also banned. The student-athlete shall be held accountable for all drugs within the banned-drug class regardless of whether they have been specifically identified. There is no complete list of banned substances.

Substances and Methods Subject to Restrictions:

- Blood and gene doping.
- Local anesthetics (permitted under some conditions).
- Manipulation of urine samples.
- Beta-2 agonists (permitted only by inhalation with prescription).
- Tampering of urine samples.

NJSIAA Nutritional/Dietary Supplements:

Warning: Before consuming any nutritional/dietary supplement product, review the product and its label with your athletics department staff!

- Nutritional/Dietary supplements, including vitamins and minerals, are not well regulated and may cause a
 positive drug test.
- Student-athletes have tested positive and lost their eligibility using nutritional/dietary supplements.
- Many nutritional/dietary supplements are contaminated with banned substances not listed on the label.
- Any product containing a nutritional/dietary supplement ingredient is taken at your own risk.

Athletics department staff should provide guidance to student-athletes about supplement use, including a directive to have any product checked by qualified staff members before consuming. The NJSIAA subscribes only to Drug Free Sport AXISTM for authoritative review of label ingredients in medications and nutritional/dietary supplements. Contact the Drug Free Sport AXIS at www.dfsaxis.com (password: njsports).

Some Examples of Substances in Each NJSIAA Banned Drug Class.

THERE IS NO COMPLETE LIST OF BANNED SUBSTANCES. DO NOT RELY ON THIS LIST TO RULE OUT ANY LABEL INGREDIENT.

Stimulants:

amphetamine (Adderall); caffeine (guarana); cocaine; ephedrine; methamphetamine; methylphenidate (Ritalin); synephrine (bitter orange); dimethylamylamine (DMAA, methylhexanamine); "bath salts" (mephedrone); Octopamine; hordenine; dimethylbutylamine (DMBA, AMP, 4-amino methylpentane citrate); phenethylamines (PEAs); dimethylhexylamine (DMHA, Octodrine); heptaminol etc. exceptions: phenylephrine and pseudoephedrine are not banned.

Anabolic Agents (sometimes listed as a chemical formula, such as 3,6,17-androstenetrione):

Androstenedione; boldenone; clenbuterol; DHEA (7-Keto); epi-trenbolone; testosterone; etiocholanolone; methasterone; methandienone; nandrolone; norandrostenedione; stanozolol; stenbolone; trenbolone; SARMS (ostarine, ligandrol, LGD-4033, S-23, RAD140)); DHCMT (oral turanibol) etc.

Alcohol and Beta Blockers:

alcohol; atenolol; metoprolol; nadolol; pindolol; propranolol; timolol; etc.

Diuretics and Masking Agents:

bumetanide; chlorothiazide; furosemide; hydrochlorothiazide; probenecid; spironolactone (canrenone); triameterene; trichlormethiazide; etc.

exceptions: finasteride is not banned

Narcotics:

Buprenorphine; dextromoramide; diamorphine (heroin); fentanyl, and its derivatives; hydrocodone; hydromorphone; methadone; morphine; nicomorphine; oxycodone; oxymorphone; pentazocine; pethidine

Cannabinoids:

marijuana; tetrahydrocannabinol (THC); synthetic cannabinoids (e.g., spice, K2, JWH-018, JWH-073)

Peptide Hormones, growth factors, related substances and mimetics

growth hormone(hGH); human chorionic gonadotropin (hCG); erythropoietin (EPO); IGF-1 (colostrum, deer antier velvet); etc.

exceptions: insulin, Synthroid are not banned

Hormone and metabolic modulators (anti-estrogens):

anastrozole; tamoxifen; formestane; ATD; SERMS (clomiphene, nolvadex); Arimidex; clomid; evista; fulvestrant; aromatase inhibitors (Androst-3,5-dien-7,17-dione), letrozole; etc.

Beta-2 Agonists:

bambuterol; formoterol; salbutamol; salmeterol; higenamine; norcoclaurine; etc.

Any substance that is chemically related to one of the above classes, even if it is not listed as an example, is also banned!

Information about ingredients in medications and nutritional/dietary supplements can be obtained by contacting Drug Free Sport AXIS, www.dfsaxis.com password njsports.

It is your responsibility to check with the appropriate or designated athletics staff before using any substance.

Sports-Related Concussion and Head Injury Fact Sheet (State of New Jersey)

A concussion is a brain injury that can be caused by a blow to the head or body. Concussions are a type of Traumatic Brain Injury (TBI), which can range from mild to severe and can disrupt the way the brain normally functions. Concussions can cause significant and sustained neuropsychological impairment affecting problem solving, planning, memory, attention, concentration, and behavior.

The Centers for Disease Control and Prevention estimates that 300,000 concussions are sustained during sports related activities nationwide, and more than 62,000 concussions are sustained each year in high school contact sports. Second-impact syndrome occurs when a person sustains a second concussion while still experiencing symptoms of a previous concussion. It can lead to severe impairment and even death of the victim.

Legislation (P.L. 2010, Chapter 94) signed on December 7, 2010, mandated measures to be taken in order to ensure the safety of K-12 student-athletes involved in interscholastic sports in New Jersey. It is imperative that athletes, coaches, and parent/guardians are educated about the nature and treatment of sports related concussions and other head injuries. The legislation states that:

- All schools that participate in interscholastic sports will distribute annually this educational fact sheet to all student athletes and obtain a signed acknowledgement from each parent/guardian and student-athlete.
- Each school district shall develop a written policy describing the prevention and treatment of sportsrelated concussion and other head injuries sustained by interscholastic student-athletes.
- Any student-athlete who participates in an interscholastic sports program and is suspected of sustaining a concussion will be immediately removed from competition or practice. The student-athlete will not be allowed to return to competition or practice until he/she has written clearance from a physician trained in concussion treatment and has completed his/her district's graduated return-to-play protocol.

Quick Facts

- · Most concussions do not involve loss of consciousness
- You can sustain a concussion even if you do not hit your head. A blow elsewhere on the body can transmit an "impulsive" force to the brain and cause a concussion

Signs of Concussions (Observed by Coach, Athletic Trainer, Parent/Guardian)

- · Appears dazed or stunned
- Forgets plays or demonstrates short term memory difficulties (e.g. unsure of game, opponent)
- Exhibits difficulties with balance, coordination, concentration, and attention
- · Answers questions slowly or inaccurately
- Demonstrates behavior or personality changes
- · Is unable to recall events prior to or after the hit or fall

Symptoms of Concussion (Reported by Student-Athlete)

- Headache
- Nausea/vomiting
- · Balance problems or dizziness
- · Double vision or changes in vision
- Sensitivity to light/sound
- Feeling of sluggishness or fogginess
- · Difficulty with concentration, short term memory, and/or confusion

OPIOID USE AND MISUSE EDICATIONAL FACTORIES

Keeping Student-Athletes Safe

School athletics can serve an integral role in students' development. In addition to providing healthy forms of exercise, school athletics foster friendships and camaraderie, promote sportsmanship and fair play, and instill the value of competition.

Unfortunately, sports activities may also lead to injury and, in rare cases, result in pain that is severe or long-lasting enough to require a prescription opioid painkiller. It is important to understand that overdoses from opioids are on the rise and are killing Americans of all ages and backgrounds. Families and communities across the country are coping with the health, emotional and economic effects of this epidemic.²

This educational fact sheet, created by the New Jersey Department of Education as required by state law (N.J.S.A. 18A:40-41.10), provides information concerning the use and misuse of opioid drugs in the event that a health care provider prescribes a student-athlete or cheerleader an opioid for a sports-related injury. Student-athletes and cheerleaders participating in an interscholastic sports program (and their parent or guardian, if the student is under age 18) must provide their school district written acknowledgment of their receipt of this fact sheet.

Hoyen Addings (digam (duning)

In some cases, student-athletes are prescribed these medications. According to research, about a third of young people studied obtained pills from their own previous prescriptions (i.e., an unfinished prescription used outside of a physician's supervision), and 83 percent of adolescents had unsupervised access to their prescription medications.³ It is important for parents to understand the possible hazard of having unsecured prescription medications in their households. Parents should also understand the importance of proper storage and disposal of medications, even if they believe their child would not engage in non-medical use or diversion of prescription medications.

What is estimated optible tree

According to the National Council on Alcoholism and Drug Dependence, 12 percent of male athletes and 8 percent of female athletes had used prescription opioids in the 12-month period studied.³ In the early stages of abuse, the athlete may exhibit unprovoked nausea and/or vomiting. However, as he or she develops a tolerance to the drug, those signs will diminish. Constipation is not uncommon, but may not be reported. One of the most significant indications of a possible opioid addiction is an athlete's decrease in academic or athletic performance, or a lack of interest in his or her sport. If these warning signs are noticed, best practices call for the student to be referred to the appropriate professional for screening.⁴ such as provided through an evidence-based practice to identify problematic use, abuse and dependence on illicit drugs (e.g., Screening, Brief Intervention, and Referral to Treatment (SBIRT)) offered through the New Jersey Department of Health.


What Are Some Ways Opioid Use and Misuse Can Be Prevented?

According to the New Jersey State Interscholastic Athletic Association (NJSIAA) Sports Medical Advisory Committee chair, John P. Kripsak, D.O., "Studies indicate that about 80 percent of heroin users started out by abusing narcotic painkillers."

The Sports Medical Advisory Committee, which includes representatives of NJSIAA member schools as well as experts in the field of healthcare and medicine, recommends the following:

- The pain from most sports-related injuries can be managed with non-narcotic medications such as acetaminophen, nonsteroidal anti-inflammatory medications like ibuprofen, naproxen or aspirin. Read the label carefully and always take the recommended dose, or follow your doctor's instructions. More is not necessarily better when taking an over-the-counter (OTC) pain medication, and it can lead to dangerous side effects.
- Ice therapy can be utilized appropriately as an anesthetic.
- Always discuss with your physician exactly what is being prescribed for pain and request to avoid narcotics.
- In extreme cases, such as severe trauma or post-surgical pain, opioid pain medication should not be prescribed for more than five days at a time;
- Parents or guardians should always control the dispensing of pain medications and keep them in a safe, non-accessible location; and
- Unused medications should be disposed of immediately upon cessation of use. Ask your pharmacist about drop-off locations or home disposal kits like Deterra or Medsaway.

According to NJSIAA Sports
Medical Advisory Committee chair,
John P. Kripsak, D.O., "Studies
Indicate that about 80 percent of
heroin users started out by abusing
narcotic painkillers."




NJ Health

STATE OF NEW JERSEY DEPARTMENT OF HEALTH

NJSIAA SPORTS MEDICAL ADVISORY COMMITTEE



Karan Chauhan Parsippany Hills High School, Permanent Student Representative New Jersey State Board of Education



Even With Proper Training and Prevention, Sports Injuries May Occur

There are two kinds of sports injuries. Acute injuries happen suddenly, such as a sprained ankle or strained back. Chronic injuries may happen after someone plays a sport or exercises over a long period of time, even when applying overuse-preventative techniques.5

Athletes should be encouraged to speak up about injuries, coaches should be supported in injury-prevention decisions, and parents and young athletes are encouraged to become better educated about sports safety.6

What Are Some Ways to Reduce the Risk of Injury?

Half of all sports medicine injuries in children and teens are from overuse. An overuse injury is damage to a bone, muscle, ligament, or tendon caused by repetitive stress without allowing time for the body to heal. Children and teens are at increased risk for overuse injuries because growing bones are less resilient to stress. Also, young athletes may not know that certain symptoms are signs of overuse.

The best way to deal with sports injuries is to keep them from happening in the first place. Here are some recommendations to consider:

PREPARE Obtain the preparticipation physical evaluation prior to participation on a school-sponsored interscholastic or intramural athletic team or squad.



CONDITIONING Maintain a good fitness level during the season and offseason. Also important are proper warm-up and cooldown

PLAY SMART Try a variety of sports and consider specializing in one sport before late adolescence to help avoid overuse injuries.



ADEQUATE HYDRATION Keep the body hydrated to help the heart more easily pump blood to muscles, which helps muscles work efficiently.

TRAINING Increase weekly training time, mileage or repetitions no more than 10 percent per week. For example, if running 10 miles one week, increase to 11 miles the following week. Athletes should also cross-train and perform sport-specific drills in different ways, such as running in a swimming pool instead of only running on the road.



REST UP Take at least one day off per week from organized activity to recover physically and mentally. Athletes should take a combined three months off per year from a specific sport (may be divided throughout the year in one-month increments). Athletes may remain physically active during rest periods through alternative low-stress activities such as stretching, yoga or walking.

PROPER EQUIPMENT Wear appropriate and properly fitted protective equipment such as pads (neck, shoulder, elbow, chest, knee, and shin), helmets, mouthpieces, face guards, protective cups, and eyewear. Do not assume that protective gear will prevent all injuries while performing more dangerous or risky activities.

Resources for Parents and Students on Preventing Substance Misuse and Abuse

The following list provides some examples of resources:

National Council on Alcoholism and Drug Dependence - NJ promotes addiction treatment and recovery.

New Jersey Department of Health, Division of Mental Health and Addiction Services is committed to providing consumers and families with a wellness and recovery-oriented model of care.

New Jersey Prevention Network includes a parent's quiz on the effects of opioids.

Operation Prevention Parent Toolkit is designed to help parents learn more about the oploid epidemic, recognize warning signs, and open lines of communication with their children and those in the community.

Parent to Parent NJ is a grassroots coalition for families and children struggling with alcohol and drug addiction.

Partnership for a Drug Free New Jersey is New Jersey's anti-drug alliance created to localize and strengthen drug-prevention media efforts to prevent unlawful drug use, especially among young people.

The Science of Addiction: The Stories of Teens shares common misconceptions about opioids through the voices of teens.

Youth IMPACTing NJ is made up of youth representatives from coalitions across the state of New Jersey who have been impacting their communities and peers by spreading the word about the dangers of underage drinking, marijuana use, and other substance misuse.

References 1 Massachusetts Technical Assistance Partnership

- for Prevention
- ² Centers for Disease Control and Prevention 3 New Jersey State Interscholastic Athletic
- Association (NJSIAA) Sports Medical Advisory Committee (SMAC)
- 4 Athletic Management, David Csillan, athletic trainer, Ewing High School, NJSIAA SMAC
- 5 National Institute of Arthritis and Musculoskelelal and Skin Diseases
- USA TODAY
- 7 American Academy of Pediatrics

An online version of this fact sheet is available on the New Jersey Department of Education's Alcohol, Tobacco, and Other Drug Use webpage. Updated Jan. 30, 2018.

SPORTS-RELATED
EYE INJURIES:

AN EDUCATIONAL FACT SHEET FOR PARENTS



Participating in sports and recreational activities is an important part of a healthy, physically active lifestyle for children. Unfortunately, injuries can, and do, occur. Children are at particular risk for sustaining a sports-related eye injury and most of these injuries can be prevented. Every year, more than 30,000 children sustain serious sports-related eye injuries. Every 13 minutes, an emergency room in the United States treats a sports-related eye injury.\(^1\) According to the National Eye Institute, the sports with the highest rate of eye injuries are: baseball/softball, ice hockey, racquet sports, and basketball, followed by fencing, lacrosse, paintball and boxing.

Thankfully, there are steps that parents can take to ensure their children's safety on the field, the court, or wherever they play or participate in sports and recreational activities.

Prevention of Sports-Related Eye Injuries

Approximately 90% of sports-related eye injuries can be prevented with simple precautions, such as using protective eyewear.² Each sport has a certain type of recommended protective eyewear, as determined by the American Society for Testing and Materials (ASTM). Protective eyewear should sit comfortably on the face. Poorly fitted equipment may be uncomfortable, and may not offer the best eye protection. Protective eyewear for sports includes, among other things, safety goggles and eye guards, and it should be made of polycarbonate lenses, a strong, shatterproof plastic. Polycarbonate lenses are much stronger than regular lenses.³

Health care providers (HCP), including family physicians, ophthalmologists, optometrists, and others, play a critical role in advising students, parents and guardians about the proper use of protective eyewear. To find out what kind of eye protection is recommended, and permitted for your child's sport, visit the National Eye Institute at http://www.nei.nih.gov/sports/findingprotection.asp. Prevent Blindness America also offers tips for choosing and buying protective eyewear at http://www.preventblindness.org/tips-buying-sports-eye-protectors, and http://www.preventblindness.org/ recommended-sports-eye-protectors.

It is recommended that all children participating in school sports or recreational sports wear protective eyewear. Parents and coaches need to make sure young athletes protect their eyes, and properly gear up for the game. Protective eyewear should be part of any uniform to help reduce the occurrence of sports-related eye injuries. Since many youth teams do not require eye protection, parents may need to ensure that their children wear safety glasses or goggles whenever they play sports. Parents can set a good example by wearing protective eyewear when they play sports.

National Eye Institute, National Eye Health Education Program, Sports-Related Eye Injuries: What You Need to Know and Tips for Prevention, www.nel.nih.gov/sports/pdf/sportsrelatedeye/njuries.pdf, December 26, 2013.

Rodriguez, Jorge O., D.O., and Lavina, Adrian M., M.D., Prevention and Treatment of Common Eye Injuries in Sports, http://www.aafp.org/afp/2003/0401/p1481.html, September 4, 2014; National Eye Health Education Program, Sports-Related Eye Injuries: What You Need to Know and Tips for Prevention, www.nei.nih.gov/sports/pdf/sportsrelatedeyeInjuries.pdf, December 26, 2013.

Bedinghaus, Troy, O.D., Sports Eye Injuries, http://vision.about.com/od/emergencyeyecare/a/Sports_Injuries.htm, December 27, 2013.

Most Common
Types of Eye
Injuries

The most common types of eye injuries that can result from sports injuries are blunt injuries, corneal abrasions and penetrating injuries.

- Blunt injuries: Blunt injuries occur when the eye is suddenly compressed by impact from an object. Blunt injuries, often caused by tennis balls, racquets, fists or elbows, sometimes cause a black eye or hyphema (bleeding in front of the eye). More serious blunt injuries often break bones near the eye, and may sometimes seriously damage important eye structures and/or lead to vision loss.
- Corneal abrasions: Corneal abrasions are painful scrapes on the outside of the eye, or the cornea. Most corneal abrasions eventually heal on their

own, but a doctor can best assess the extent of the abrasion, and may prescribe medication to help control the pain. The most common cause of a sports-related corneal abrasion is being poked in the eye by a finger.

- ◆ Penetrating injuries: Penetrating injuries are caused by a foreign object piercing the eye. Penetrating injuries are very serious, and often result in severe damage to the eye. These injuries often occur when eyeglasses break while they are being worn. Penetrating injuries must be treated quickly in order to preserve vision.⁴
- Pain when looking up and/or down, or difficulty seeing;
- Tenderness;
- Sunken eye;
- Double vision;
- Severe eyelid and facial swelling;
- Difficulty tracking;

Signs or Symptoms of an Eye Injury



- The eye has an unusual pupil size or shape;
- Blood in the clear part of the eye;
- Numbness of the upper cheek and gum; and/or
- Severe redness around the white part of the eye.

What to do if a
Sports-Related
Eye Injury
Occurs

If a child sustains an eye injury, it is recommended that he/she receive immediate treatment from a licensed HCP (e.g., eye doctor) to reduce the risk of serious damage, including blindness. It is also recommended that the child, along with his/her parent or guardian, seek guidance from the HCP regarding the appropriate amount of time to wait before returning to sports competition or practice after sustaining an eye injury. The school nurse and the child's teachers should also be notified when a child sustains an eye injury. A parent or guardian should also provide the school nurse with a physician's note detailing the nature of the eye injury, any diagnosis, medical orders for

the return to school, as well as any prescription(s) and/or treatment(s) necessary to promote healing, and the safe resumption of normal activities, including sports and recreational activities.

According to the American Family Physician Journal, there are several guidelines that should be followed when students return to play after sustaining an eye injury. For

Return to Play and Sports example, students who have sustained significant ocular injury should receive a full examination and clearance by an ophthalmologist or optometrist. In addition, students should not return to play until the period of time recommended by their HCP has elapsed. For more minor eye injuries, the athletic trainer may determine that

it is safe for a student to resume play based on the nature of the injury, and how the student feels. No matter what degree of eye injury is sustained, it is recommended that students wear protective eyewear when returning to play and immediately report any concerns with their vision to their coach and/or the athletic trainer.

Additional information on eye safety can be found at http://isee.nei.nih.gov and http://www.nei.nih.gov/sports.

Website Resources

- Sudden Death in Athletes http://tinyurl.com/m2gjmvq
- Hypertrophic Cardiomyopathy Association www.4hcm.org
- American Heart Association www.heart.org

Collaborating Agencies:

American Academy of Pediatrics **New Jersey Chapter** 3836 Quakerbridge Road, Suite 108 Hamilton, NJ 08619 (p) 609-842-0014 (f) 609-842-0015



American Heart Association


1 Union Street, Suite 301 Robbinsville, NJ, 08691 (p) 609-208-0020 www.heart.org

www.aapnj.org



New Jersey Department of Education PO Box 500

Trenton, NJ 08625-0500 (p) 609-292-5935 www.state.nj.us/education/




P.O. Box 360 Trenton, NJ 08625-0360 (p) 609-292-7837 www.state.ni.us/health

Lead Author: American Academy of Pediatrics, **New Jersey Chapter**

Written by: Initial draft by Sushma Raman Hebbar, MD & Stephen G. Rice, MD PhD

Additional Reviewers: NJ Department of Education, NJ Department of Health and Senior Services, American Heart Association/New Jersey Chapter, NJ Academy of Family Practice, Pediatric Cardiologists, New Jersey State School Nurses


Revised 2014: Nancy Curry, EdM; Christene DeWitt-Parker, MSN, CSN, RN; Lakota Kruse, MD, MPH; Susan Martz, EdM; Stephen G. Rice, MD; Jeffrey Rosenberg, MD, Louis Teichholz, MD; Perry Weinstock, MD



American Academy of Pediatrics DEDICATED TO THE HEALTH OF ALL CHILDREN-



American Heart Learn and Live



udden death in young athletes between the ages of 10 and 19 is very rare. What, if anything, can be done to prevent this kind of tragedy?

What is sudden cardiac death in the young athlete?

Sudden cardiac death is the result of an unexpected failure of proper heart function, usually (about 60% of the time) during or immediately after exercise without trauma. Since the heart stops pumping adequately, the athlete quickly collapses, loses consciousness, and ultimately dies unless normal heart rhythm is restored using an automated external defibrillator (AED).

How common is sudden death in young athletes?

Sudden cardiac death in young athletes is a very rare. About 100 such deaths are reported in the United States per year. The chance of sudden death occurring to any individual high school athlete is about one in 200,000 per year.

Sudden cardiac death is more common: in males than in females; in football and basketball than in other sports; and in African-Americans than in other races and ethnic groups.

What are the most common causes?

Research suggests that the main cause is a loss of proper heart rhythm, causing the heart to guiver instead of pumping blood to the brain and body. This is called ventricular fibrillation (ven-TRICK-you-lar fibroo-LAY-shun). The problem is usually caused by one of several cardiovascular abnormalities and electrical diseases of the heart that go unnoticed in healthy-appearing athletes.

The most common cause of sudden death in an athlete is hypertrophic cardiomyopathy (hi-per-TRO-fic CAR- dee-oh-my-OP-a-thee) also called HCM. HCM is a disease of the heart, with abnormal thickening of the heart muscle, which can cause serious heart rhythm problems and blockages to blood flow. This genetic disease runs in families and usually develops gradually over many years.

The second most likely cause is congenital (con-JEN-it-al) (i.e., present from birth) abnormalities of the coronary

> arteries. This means that these blood vessels are connected to the main blood vessel of the heart in an abnormal way. This differs from blockages that may occur when people get older (commonly called "coronary artery disease," which may lead to a heart attack).

SUDDEN CARDIAC DEATH IN YOUNG ATHLETES

Other diseases of the heart that can lead to will sudden death in young people include:

- Myocarditis (my-oh-car-DIE-tis), an acute inflammation of the heart muscle (usually due to a virus).
- Dilated cardiomyopathy, an enlargement of the heart for unknown reasons.
- Long QT syndrome and other electrical abnormalities of the heart which cause abnormal fast heart rhythms that can also run in families.
- Marfan syndrome, an inherited disorder that affects heart valves, walls of major arteries, eyes and the skeleton. It is generally seen in unusually tall athletes, especially if being tall is not common in other family members.

Are there warning signs to watch for?

In more than a third of these sudden cardiac deaths, there were warning signs that were not reported or taken seriously. Warning signs are:

- Fainting, a seizure or convulsions during physical activity;
- Fainting or a seizure from emotional excitement, emotional distress or being startled;
- Dizziness or lightheadedness, especially during exertion;
- Chest pains, at rest or during exertion;
- Palpitations awareness of the heart beating unusually (skipping, irregular or extra beats) during athletics or during cool down periods after athletic participation;
- · Fatigue or tiring more quickly than peers; or
- Being unable to keep up with friends due to shortness of breath (labored breathing).

What are the current recommendations for screening young athletes?

New Jersey requires all school athletes to be examined by their primary care physician ("medical home") or school physician at least once per year. The New Jersey Department of Education requires use of the specific Preparticipation Physical Examination Form (PPE).

This process begins with the parents and student-athletes answering questions about symptoms during exercise (such as chest pain, dizziness, fainting, palpitations or shortness of breath); and questions about family health history.

The primary healthcare provider needs to know if any family member died suddenly during physical activity or during a seizure. They also need to know if anyone in the family under the age of 50 had an unexplained sudden death such as drowning or car accidents. This information must be provided annually for each exam because it is so essential to identify those at risk for sudden cardiac death.

The required physical exam includes measurement of blood pressure and a careful listening examination of the heart, especially for murmurs and rhythm abnormalities. If there are no warning signs reported on the health history and no abnormalities discovered on exam, no further evaluation or testing is recommended.

Are there options privately available to screen for cardiac conditions?

Technology-based screening programs including a 12-lead electrocardiogram (ECG) and echocardiogram (ECHO) are noninvasive and painless options parents may consider in addition to the required

expensive and are not currently advised by the American Academy of Pediatrics and the American College of Cardiology unless the PPE reveals an indication for these tests. In addition to the expense, other limitations of technology-based tests include the possibility of "false positives" which leads to unnecessary stress for the student and parent or guardian as well as unnecessary restriction from athletic participation.

The United States Department of Health and Human Services offers risk assessment options under the Surgeon General's Family History Initiative available at http://www.hhs.gov/familyhistory/index.html.

When should a student athlete see a heart specialist?

If the primary healthcare provider or school physician has concerns, a referral to a child heart specialist, a pediatric cardiologist, is recommended. This specialist will perform a more thorough evaluation, including an electrocardiogram (ECG), which is a graph of the electrical activity of the heart. An echocardiogram, which is an ultrasound test to allow for direct visualization of the heart structure, will likely also be done. The specialist may also order a treadmill exercise test and a monitor to enable a longer recording of the heart rhythm. None of the testing is invasive or uncomfortable.

Can sudden cardiac death be prevented just through proper screening?

A proper evaluation should find most, but not all, conditions that would cause sudden death in the athlete. This is because some diseases are difficult to uncover and may only develop later in life. Others can develop following a

normal screening evaluation, such as an infection of the heart muscle from a virus.

This is why screening evaluations and a review of the family health history need to be performed on a yearly basis by the athlete's primary healthcare provider. With proper screening and evaluation, most cases can be identified and prevented.

Why have an AED on site during sporting events?

The only effective treatment for ventricular fibrillation is immediate use of an automated external defibrillator (AED). An AED can restore the heart back into a normal rhythm. An AED is also life-saving for ventricular fibrillation caused by a blow to the chest overthe heart (commotio cordis).

N.J.S.A. 18A:40-41a through c, known as "Janet's Law," requires that at any schoolsponsored athletic event or team practice in. New Jersey public and nonpublic schools including any of grades K through 12, the following must be available:

- An AED in an unlocked location on school property within a reasonable proximity to the athletic field or gymnasium; and
- A team coach, licensed athletic trainer, or other designated staff member if there is no coach or licensed athletic trainer present, certified in cardiopulmonary resuscitation (CPR) and the use of the AED; or
- A State-certified emergency services provider or other certified first responder.

The American Academy of Pediatrics recommends the AED should be placed in central location that is accessible and ideally no more than a 1 to 1¹/₂ minute walk from any location and that a call is made to activate 911 emergency system while the AED is being retrieved.